


THIRD EDITION

Strength Training Anatomy



OVER
1 MILLION
COPIES
SOLD!

Frédéric Delavier

Strength Training Anatomy

Over 1 million copies sold! With new exercises, additional stretches, and more of Frédéric Delavier's signature illustrations, you'll gain a whole new understanding of how muscles perform during strength exercises. This one-of-a-kind best-seller combines the visual detail of top anatomy texts with the best of strength training advice.

Many books explain what muscles are used during exercise, but no other resource brings the anatomy to life like *Strength Training Anatomy*. Over 600 full-color illustrations reveal the primary muscles worked along with all the relevant surrounding structures, including bones, ligaments, tendons, and connective tissue.

Like having an X-ray for each exercise, the anatomical depictions show both superficial and deep layers and detail how various setup positions affect muscle recruitment and emphasize underlying structures. New pages show common strength training injuries in a fascinating light and offer precautions to help you exercise safely.

Author and illustrator **Frédéric Delavier** is the former editor in chief of the French publication *PowerMag*. He is a journalist for *Le Monde du Muscle* and a contributor to *Men's Health Germany* and several other strength training publications.



Human Kinetics

ISBN-13: 978-0-7360-9226-5

ISBN-10: 0-7360-9226-9



5 1 9 9 5



9 780736 092265


US \$19.95
CAN \$21.95

frédéric delavier

STRENGTH TRAINING ANATOMY

THIRD EDITION

常州大学图书馆
藏书章



HUMAN KINETICS

For my father

Library of Congress Cataloging-in-Publication Data

Delavier, Frédéric.

[Guide des mouvements de musculation. English]

Strength training anatomy / Frédéric Delavier. -- 3rd ed.

p. cm.

ISBN-13: 978-0-7360-9226-5 (soft cover)

ISBN-10: 0-7360-9226-9 (soft cover)

1. Muscles--Anatomy. 2. Weight training. 3. Muscle strength. I. Title.

QM151.D454 2010

611'.73--dc22

2009045227

ISBN-10: 0-7360-9226-9 (print)

ISBN-13: 978-0-7360-9226-5 (print)

Copyright © 2010, 2006, 2001 by Éditions Vigot, 23 rue de l'École de Médecine, 75006 Paris, France

All rights reserved. Except for use in a review, the reproduction or utilization of this work in any form or by any electronic, mechanical, or other means, now known or hereafter invented, including xerography, photocopying, and recording, and in any information storage and retrieval system, is forbidden without the written permission of the publisher.

This publication is written and published to provide accurate and authoritative information relevant to the subject matter presented. It is published and sold with the understanding that the author and publisher are not engaged in rendering legal, medical, or other professional services by reason of their authorship or publication of this work. If medical or other expert assistance is required, the services of a competent professional person should be sought.

This book is a revised edition of *Guide des Mouvements de Musculation*, 5th edition, published in 2010 by Éditions Vigot.

Illustrator: Frédéric Delavier

Human Kinetics books are available at special discounts for bulk purchase. Special editions can also be created to specification. For details, contact the Special Sales Manager at Human Kinetics.

Printed in France - L57439A 10 9 8 7 6 5

Human Kinetics

Web site: www.HumanKinetics.com

United States: Human Kinetics

P.O. Box 5076

Champaign, IL 61825-5076

800-747-4457

e-mail: humank@hkusa.com

Canada: Human Kinetics

475 Devonshire Road Unit 100

Windsor, ON N8Y 2L5

800-465-7301 (in Canada only)

e-mail: info@hkcanada.com

Europe: Human Kinetics

107 Bradford Road

Stanningley

Leeds LS28 6AT, United Kingdom

+44 (0) 113 255 5665

e-mail: hk@hkeurope.com

Australia: Human Kinetics

57A Price Avenue

Lower Mitcham, South Australia 5062

08 8372 0999

e-mail: info@hkaustralia.com

New Zealand: Human Kinetics

P.O. Box 80

Torrens Park, South Australia 5062

0800 222 062

e-mail: info@hknewzealand.com

CONTENTS

| | | |
|---|----------------|-----|
| 1 | ARMS..... | 4 |
| 2 | SHOULDERS..... | 32 |
| 3 | CHEST..... | 61 |
| 4 | BACK..... | 85 |
| 5 | LEGS..... | 122 |
| 6 | BUTTOCKS..... | 155 |
| 7 | ABDOMEN..... | 170 |



frédéric delavier

STRENGTH TRAINING ANATOMY

THIRD EDITION



HUMAN KINETICS

For my father

Library of Congress Cataloging-in-Publication Data

Delavier, Frédéric.

[Guide des mouvements de musculation. English]

Strength training anatomy / Frédéric Delavier. -- 3rd ed.

p. cm.

ISBN-13: 978-0-7360-9226-5 (soft cover)

ISBN-10: 0-7360-9226-9 (soft cover)

1. Muscles--Anatomy. 2. Weight training. 3. Muscle strength. I. Title.

QM151.D454 2010

611'.73--dc22

2009045227

ISBN-10: 0-7360-9226-9 (print)

ISBN-13: 978-0-7360-9226-5 (print)

Copyright © 2010, 2006, 2001 by Éditions Vigot, 23 rue de l'École de Médecine, 75006 Paris, France

All rights reserved. Except for use in a review, the reproduction or utilization of this work in any form or by any electronic, mechanical, or other means, now known or hereafter invented, including xerography, photocopying, and recording, and in any information storage and retrieval system, is forbidden without the written permission of the publisher.

This publication is written and published to provide accurate and authoritative information relevant to the subject matter presented. It is published and sold with the understanding that the author and publisher are not engaged in rendering legal, medical, or other professional services by reason of their authorship or publication of this work. If medical or other expert assistance is required, the services of a competent professional person should be sought.

This book is a revised edition of *Guide des Mouvements de Musculation*, 5th edition, published in 2010 by Éditions Vigot.

Illustrator: Frédéric Delavier

Human Kinetics books are available at special discounts for bulk purchase. Special editions can also be created to specification. For details, contact the Special Sales Manager at Human Kinetics.

Printed in France - L57439A 10 9 8 7 6 5

Human Kinetics

Web site: www.HumanKinetics.com

United States: Human Kinetics

P.O. Box 5076

Champaign, IL 61825-5076

800-747-4457

e-mail: humank@hkusa.com

Canada: Human Kinetics

475 Devonshire Road Unit 100

Windsor, ON N8Y 2L5

800-465-7301 (in Canada only)

e-mail: info@hkcanada.com

Europe: Human Kinetics

107 Bradford Road

Stanningley

Leeds LS28 6AT, United Kingdom

+44 (0) 113 255 5665

e-mail: hk@hkeurope.com

Australia: Human Kinetics

57A Price Avenue

Lower Mitcham, South Australia 5062

08 8372 0999

e-mail: info@hkaustralia.com

New Zealand: Human Kinetics

P.O. Box 80

Torrens Park, South Australia 5062

0800 222 062

e-mail: info@hknewzealand.com




CONTENTS





| | | |
|---|----------------|-----|
| 1 | ARMS..... | 4 |
| 2 | SHOULDERS..... | 32 |
| 3 | CHEST..... | 61 |
| 4 | BACK..... | 85 |
| 5 | LEGS..... | 122 |
| 6 | BUTTOCKS..... | 155 |
| 7 | ABDOMEN..... | 170 |



1



ARMS

| | |
|---|-----------|
| 1 DUMBBELL CURLS | 6 |
| 2 CONCENTRATION CURLS | 7 |
| 3 INCLINE DUMBBELL CURLS | 8 |
| 4 HAMMER CURLS | 9 |
| 5 LOW-PULLEY CURLS | 10 |
| 6 HIGH-PULLEY CURLS | 11 |
| 7 BARBELL CURLS | 12 |
|  ELBOW STRUCTURE AND ITS EFFECT ON TRAINING | 13 |
| 8 MACHINE CURLS | 14 |
| 9 PREACHER CURLS | 15 |
| 10 REVERSE WRIST CURLS | 16 |
| 11 WRIST CURLS | 17 |
| 12 REVERSE BARBELL CURLS | 18 |
|  STRETCHING THE FOREARM MUSCLES | 19 |
| 13 PUSH-DOWNS | 20 |
| 14 REVERSE PUSH-DOWNS | 21 |
| 15 ONE-ARM REVERSE PUSH-DOWNS | 22 |
| 16 OVERHEAD CABLE TRICEPS EXTENSIONS | 23 |
| 17 LYING TRICEPS EXTENSIONS | 24 |
| 18 LYING DUMBBELL TRICEPS EXTENSIONS | 25 |
| 19 ONE-ARM OVERHEAD DUMBBELL TRICEPS EXTENSIONS | 26 |
| 20 TRICEPS KICKBACKS | 27 |
| 21 SEATED DUMBBELL TRICEPS EXTENSIONS | 28 |
| 22 SEATED E-Z BAR TRICEPS EXTENSIONS | 29 |
| 23 TRICEPS DIPS | 30 |
|  STRETCHING THE TRICEPS | 31 |



1

DUMBBELL CURLS




Comment: This exercise takes the biceps through its complete range of motion, which includes flexion, protraction, and supination.

Sit holding a dumbbell in each hand with arms hanging down and the palms of the hands facing the body:

- Inhale and bend the elbow, rotating the palm up before the forearm reaches horizontal.
- Continue by raising the elbow at the end of the movement.

This exercise primarily uses the brachioradialis (long supinator), brachialis, biceps brachii, and anterior deltoid, and, to a lesser extent, the coracobrachialis and clavicular head of the pectoralis major.






THREE WAYS TO EXECUTE CURLS:

- 1 Emphasize biceps
- 2 Work brachioradialis intensely
- 3 Work mainly biceps and brachialis



CONCENTRATION CURLS

2



FINAL POSITION

BRACHIALIS MUSCLE



Sit holding a dumbbell with the palm facing forward and the elbow positioned against the inner thigh:



- Inhale and lift the forearm by bending the elbow.
- Exhale at the end of the effort.

This isolation exercise allows you to control the range of motion, speed, and form of the movement.

It mainly works the biceps brachii and brachialis.

3

INCLINE DUMBBELL CURLS



■ Recruited part of muscle
■ Most recruited part of muscle

Sit on an incline bench with your back leaning against the support. In each hand, hold a dumbbell with a relaxed overhand grip (thumbs face to the inside):

- Inhale and bend your elbows while externally rotating your wrists before your forearms reach horizontal so that at the end of the movement your hands are in an underhand grip (thumbs face to the outside).
- Exhale at the end of the movement.


This exercise targets the long head of the biceps (the lateral part of the muscle), which is stretched at the beginning of flexion of the elbow. This movement also works the brachioradialis and the brachialis.

Variations: Use alternating forearm curls. Increase the intensity of the effort by initiating the movement in an underhand grip.




Attention: Adjust the angle of the bench according to individual variations in shoulder flexibility. If the arm is too far back, the long head of the biceps will create excessive friction in the bicipital groove of the humerus and will strain the tendon.

INITIAL POSITION





HAMMER CURLS

4



THE MOVEMENT





Stand or sit gripping a dumbbell in each hand with the palms facing each other:

- Inhale and raise the forearms together or alternately.
- Exhale at the end of the movement.

This is the best exercise for developing the brachioradialis. It also develops the biceps brachii, brachialis, and, to a lesser degree, the extensor carpi radialis brevis and longus.

BRACHIORADIALIS MUSCLE

DISTRIBUTION OF THE MUSCLE FIBERS
ACCORDING TO THEIR SPECIFIC ACTIONS

- Deep muscle fibers in the most direct path: most frequently recruited.
- Superficial muscle fibers: recruited as intensity of effort increases.

THE LAW OF LEAST EFFORT


To minimize effort when performing an action, the muscle will first recruit the fibers in the most direct path (that is, the most linear path located deep in the muscle).

A common belief is that the greater the force, the more the deep part of the muscle is worked. But in reality, the more the intensity of effort increases, the more the superficial muscles will be recruited to perform the movement.

Furthermore, the deep and linear muscles are generally slower to contract but more resistant to repetitive movements than the lateral, more curved, and longer muscles.

5

LOW-PULLEY CURLS



VARIATION
TWO-HANDED LOW-PULLEY CURLS




THE MOVEMENT

Stand facing the machine. Grasp the handle with an underhand grip (palm facing up):

- Inhale and bend the elbow to raise the forearm.
- Exhale at the end of the movement.

This exercise focuses the effort on the biceps brachii and works the muscle intensely.

MONOARTICULAR MUSCLE AND POLYARTICULAR MUSCLE



BRACHIALIS MUSCLE

The brachialis muscle crosses at only one articulation, the elbow. It is said to be monoarticular. Its simple action mobilizes this joint; it flexes only the forearm.






BICEPS BRACHII MUSCLE

The biceps brachii muscle crosses over at more than one articulation, that of the elbow and shoulder. It is said to be polyarticular. That is, it mobilizes more than one joint, and its action is complex. The biceps brachii can bend the elbow, raise the elbow, bring the arm to the thorax, and place the forearm into supination (underhand grip).



HIGH-PULLEY CURLS

6



7

BARBELL CURLS



Stand with the back straight. Grasp the barbell with an underhand grip and hands slightly wider than shoulder-width apart:

- Inhale and raise the barbell by bending the elbows, taking care to stabilize the torso and spine by isometrically contracting the gluteal muscles, abdominal muscles, and spinal muscles.
- Exhale at the end of the movement.

This exercise mainly contracts the biceps brachii, brachialis, and, to a lesser degree, the brachioradialis, pronator teres, and the wrist flexor group.


Variations: Vary the width of the grip to work different parts of the muscle more intensely:

- Placing the hands farther apart isolates the short head of the biceps brachii.
- Placing the hands closer together isolates the long head of the biceps brachii.

Raising both elbows after they are flexed increases the contraction of the biceps brachii and contracts the anterior deltoid.

To make the exercise more difficult, perform the movement with the back against a wall so that the shoulder blades don't move.

You can lift more weight and gain strength by leaning the torso back while lifting the bar; however, to prevent injury, this requires good technique and well-developed abdominal and lumbar muscles.



BARBELL CURLS



1 Narrow grip

Mainly works the long head of the biceps brachii


2 Wide grip

Mainly works the short head of the biceps brachii

ELBOW STRUCTURE AND ITS EFFECT ON TRAINING



Biceps training with an E-Z bar eases excessive wrist tension.



When training the biceps brachii using a barbell, take into account variations in each person's physical structure.


In the anatomical position (arms hanging alongside the body, palms facing forward, and thumbs pointing laterally), the angle at the elbow between the upper arm and

the forearm varies from person to person. Someone whose forearm hangs distinctly away from the body in a valgus position must break excessively at the wrist when performing a curl with a straight bar, which is painful. Therefore, these people should work with an E-Z bar to spare their wrists.

Comment: Valgus of the elbow is usually more pronounced in women.

8

MACHINE CURLS



Sit at the machine and grasp the bar with an underhand grip, arms extended and resting on the support:

- Inhale and raise the forearms.
- Exhale at the end of the movement.

This is one of the best exercises for working the biceps brachii. Fixing the arms against the support makes it impossible to "cheat."

At the beginning, the muscle tension is intense, so be sure to warm up properly using light weights. To avoid the risk of tendonitis, do not completely extend the arm.


This movement also works the brachialis and, to a lesser extent, the brachioradialis and pronator teres.

VARIATION

Performing the curl with an Atlas pulley is a great way to pump up the muscle.

PREACHER CURLS

9





Sit or stand with the arms resting on the support pad and grasp the bar with an underhand grip:

- Inhale and raise the forearms by bending the elbows. Exhale at the end of the effort.

This is one of the best exercises for isolating the biceps.



Attention: The angle of the support pad places significant tension on the forearms when the arm is completely extended. Therefore, warm up the muscles properly and begin with lighter weights.




10


REVERSE WRIST CURLS



FINAL POSITION



WRIST EXTENSORS




Sit with the forearms resting on the thighs or on a bench and grasp the bar with an overhand grip and keep the wrists relaxed:

- Inhale and raise the hands by extending at the wrists.

This exercise contracts the extensor carpi radialis longus and brevis, extensor digitorum, extensor digiti minimi, and the extensor carpi ulnaris.



Comment: This exercise strengthens the wrists, which are often vulnerable because of weak wrist extensors.

EXTENSOR MUSCLES AT THE WRIST




WRIST CURLS

11



WRIST FLEXORS




Sit with the forearms resting on the thighs or on a bench and grasp the bar with an underhand grip with wrists relaxed:

- Inhale and raise the hands by flexing at the wrists.

This exercise contracts the flexor carpi radialis, palmaris longus, flexor carpi ulnaris, and the flexors digitorum superficialis and profundus.


The latter two muscles, although located deep in the wrist, make up most of the muscle mass of the wrist flexors.




THE MOVEMENT

12


REVERSE BARBELL CURLS



MUSCLES OF THE FOREARM (LATERAL VIEW)



THE MOVEMENT



Stand with the legs slightly apart and arms extended and grasp the bar with an overhand grip (with the thumbs facing each other):


- Inhale and raise the forearms by bending the elbows.
- Exhale at the end of the movement.

This exercise works the extensor muscles of the wrist: extensor carpi radialis longus, extensor carpi radialis brevis, extensor digitorum, extensor digiti minimi, and extensor carpi ulnaris.

It also acts on the brachioradialis, brachialis, and, to a lesser degree, the biceps brachii.

Comment: This is an excellent exercise for strengthening the wrist, which is often weak because of an imbalance caused by using the wrist flexors rather than the wrist extensors. For this reason, many boxers include it in their training. Many bench press champions use it to keep their wrists from trembling under extreme weights.


STRETCHING THE FOREARM MUSCLES



STRETCHING THE WRIST EXTENSOR MUSCLES

Extend your arm straight forward and flex the wrist. Grasp the hand with your other hand and pull slowly as if trying to touch your anterior forearm with your palm while extending at the elbow.

This exercise mainly stretches the extensor carpi radialis longus and brevis, extensor digitorum, extensor digiti minimi, extensor carpi ulnaris, and anconeus.




STRETCHING THE WRIST FLEXOR MUSCLES

Extend your arm in front of you with the palm of that hand facing outward. Grasp that hand with your other hand and gently pull as if you are trying to bring the top of your hand toward you while pushing the palm out. This exercise mainly stretches the palmaris longus, flexor carpi radialis, flexor carpi ulnaris, superficial and deep finger flexors, and supinator.

13

PUSH-DOWNS



Stand facing the machine and grasp the handle with an overhand grip, keeping the elbows tucked into the body:

- Inhale and extend the forearms, keeping the elbows tucked into the body.
- Exhale at the end of the movement.

Comments: This exercise isolates the triceps and the anconeus.


The variation using a rope rather than a handle engages the lateral head of the triceps more intensely.

Performing the movement with an underhand grip requires more contribution from the medial head of the triceps.

Hold an isometric contraction for one or two seconds at the end of the movement to feel the effort more intensely.

When using heavy weights, lean forward with the torso.

Beginners can use this exercise to develop enough strength to move on to more difficult exercises.




VARIATION WITH BACK TO THE MACHINE
TO ISOLATE THE LONG HEAD OF THE TRICEPS



REVERSE PUSH-DOWNS

14




Stand facing the machine with the arms next to the body and elbows bent and grasp the handle with an underhand grip:

- Inhale and extend the forearms by straightening the elbows, keeping them tucked into the body.
- Exhale at the end of the movement.

The underhand grip isolates the medial head of the triceps brachii and precludes working with heavy weights.


When extending the forearms, the anconeus and wrist extensors also contract.

The extensor carpi ulnaris, extensor digitorum, extensor digiti minimi, and extensors carpi radialis longus and brevis keep the wrist straight with isometric contraction during the exercise.



15


ONE-ARM REVERSE PUSH-DOWNS



Stand facing the machine and grasp the handle with an underhand grip:


- Inhale and extend the forearm.
- Exhale at the end of the movement.

This exercise mainly works the lateral head of the triceps.



OVERHEAD CABLE TRICEPS EXTENSIONS

16




Sit with your back against a pulley machine. Grasp the pulley with an overhand grip, arms slightly above horizontal and elbows bent:

- Inhale and extend your forearms, taking care not to spread your elbows too far apart.
- Exhale at the end of the movement.

This exercise works the triceps brachii, anconeus, and serratus anterior, which maintains the scapulae fixed against the rib cage during the entire execution of the movement.



Notice that the elevated position of the elbows stretches the long heads of the triceps at the beginning of the movement, assisting its contraction during the effort.




Comment: Performing this exercise at an Atlas pulley allows you to more easily reproduce forearm extension with a bar while lying on a bench (see page 24).

17

LYING TRICEPS EXTENSIONS



1 Lowering the bar to the forehead focuses the work on the medial and lateral heads of the triceps brachii.




2 Lowering the bar behind the head focuses the work on the long head of the triceps brachii.

Lie on a horizontal bench and grasp the barbell with an overhand grip and the arms vertical:

- Inhale and lower the barbell to the forehead or behind the head by bending the elbows.
- Return to the initial position.
- Exhale at the end of the effort.

Comments: Because of individual variations in shoulder width, valgus angle at the elbows, and wrist flexibility, the hands can be closer or farther apart on the bar and the elbow angle more or less open during the exercise. Using an E-Z bar helps prevent excessive strain at the wrists.




VARIATION ON A MACHINE

Performing this exercise at an Atlas triceps pulley simulates the movement with a barbell and enables you to isolate the long head of the triceps brachii.

LYING DUMBBELL TRICEPS EXTENSIONS



18



Lie on a flat bench and grasp a dumbbell in each hand with the arms vertical:


- Inhale and lower the forearms by bending the elbow with a controlled movement.
- Return to the initial position.
- Exhale at the end of the effort.

This exercise works all three heads of the triceps brachii equally.



19

ONE-ARM OVERHEAD DUMBBELL TRICEPS EXTENSIONS




Sit or stand and grip a dumbbell in one hand with the arm vertical:

- Inhale and bend the elbow to lower the dumbbell behind the head to the neck.
- Return to the initial position.
- Exhale at the end of the movement.

The vertical position of the arm stretches the long head of the triceps brachii, emphasizing its contraction while working.



Comment: Contract the abdominal core to prevent arching the low back. If possible use a bench with support for the low back.



THE MOVEMENT

TRICEPS KICKBACKS

20



Stand with the knees slightly bent and lean forward at the waist, maintaining a straight back. Bend the elbow and hold the upper arm horizontally alongside the body:

- Inhale and extend the forearm.
- Exhale at the end of the movement.


This is an excellent exercise for pumping the triceps group. Perform this exercise until you feel a burn for best results.



INITIAL POSITION

21

SEATED DUMBBELL TRICEPS EXTENSIONS



Sit and grasp a dumbbell, holding it behind the neck:

- Inhale and extend the forearms.
- Exhale at the end of the movement.


The vertical position of the arms strongly stretches the long head of the triceps brachii, emphasizing its contraction while working.

Contract the abdominal core to prevent arching the low back. If possible use a bench with support for the low back.

Comment: The fibers of the three heads of the triceps attach to a common tendinous plate, which in turn is attached to the olecranon by a tendon. When the triceps contracts, the tendinous plate is drawn into the fibers of the heads of the triceps, much as a wooden plate drawn into butter. The contracted muscle bulges out from the plate, creating this characteristic horseshoe shape.

SEATED E-Z BAR TRICEPS EXTENSIONS

22



Sit or stand and grasp an E-Z bar with an overhand grip and arms vertical:


- Inhale and bend the elbows to lower the bar behind the head.
- Return to the initial position.
- Exhale at the end of the extension.

The vertical position of the arms strongly stretches the long head of the triceps brachii, emphasizing its contraction while working.

An overhand grip isolates the lateral head of the triceps brachii.




Contract the abdominal muscles and avoid arching the low back. If possible use a bench with support for the low back.

THE MOVEMENT



23

TRICEPS DIPS



Suspend the body between two benches by placing the hands on the edge of one bench and the feet on the edge of the other bench:

- Inhale, then dip by bending the elbows and rise by extending the forearms.
- Exhale at the end of the movement.

This exercise works the triceps and pectorals as well as the anterior deltoid.

Resting weights on top of the thighs increases the difficulty and intensity of the dip.

STRETCHING THE TRICEPS





Stand or sit with a very straight back. Hold one arm vertical against the side of your head and bend the elbow to 90 degrees.

- Try to bring your elbow behind your head.
- Hold the stretch for a few seconds while breathing slowly.

This exercise mainly stretches the triceps, teres major, and latissimus dorsi.

Variation: To accentuate the stretch on the triceps, perform this exercise with the elbow flexed and the opposing hand slowly pulling the elbow down behind the head.



TRICEPS STRETCH


Stand or sit with your back straight and one arm raised vertically beside your head. Bend the arm at the elbow and touch the top of the back with your hand. With the other hand grasp the elbow and slowly try to pull it behind your head. This stretches the teres major, the long head of the triceps brachii, and, to a lesser extent, the latissimus dorsi.

Variation: Pull the hand rather than the elbow. For greater intensity, place the raised arm against a wall.

Comment: These stretching exercises for the triceps help avoid tears, which may occur with heavy lifting on the triceps, but also when executing heavy pullovers or heavy vertical pulls. Vertical pulls have a tendency to pull excessively on the long head of the triceps.

2


SHOULDERS



| | |
|--|-----------|
| 1 BACK PRESSES | 33 |
| 2 SEATED FRONT PRESSES | 34 |
| 3 SEATED DUMBBELL PRESSES | 35 |
| + SHOULDER INJURIES | 36 |
| + LYING DUMBBELL PRESSES | 38 |
| HOW THE OSSEOUS MORPHOLOGY AFFECTS THE BACK PRESS | 39 |
| 4 ARNOLD PRESSES | 40 |
| 5 BENT-OVER LATERAL RAISES | 41 |
| 6 LATERAL DUMBBELL RAISES | 42 |
| 7 ALTERNATE FRONT ARM RAISES | 44 |
| 8 SIDE-LYING LATERAL RAISES | 45 |
| 9 LOW-PULLEY FRONT RAISES, OVERHAND GRIP | 46 |
| 10 LOW-PULLEY FRONT RAISES, NEUTRAL GRIP | 47 |
| STRETCHING THE ANTERIOR DELTOIDS | 48 |
| 11 HIGH-PULLEY LATERAL EXTENSIONS | 49 |
| 12 EXTERNAL ARM ROTATIONS AT A PULLEY | 50 |
| 13 LOW-PULLEY BENT-OVER LATERAL RAISES | 51 |
| 14 LOW-PULLEY LATERAL RAISES | 52 |
| IMPORTANCE OF PULLING THE SHOULDERS BACK | 53 |
| 15 ONE-DUMBBELL FRONT RAISES | 54 |
| 16 BARBELL FRONT RAISES | 55 |
| 17 UPRIGHT ROWS | 56 |
| 18 MACHINE LATERAL RAISES | 57 |
| 19 PEC DECK REAR-DELT LATERALS | 58 |
| STRETCHING THE POSTERIOR ROTATOR CUFF MUSCLES | 59 |
| STRETCHING THE SHOULDER | 60 |

BACK PRESSES

1




Sit with the back straight and hold the bar across the back of the neck with an overhand grip:

- Inhale and extend the bar straight up, keeping the low back as straight as possible.
- Exhale at the end of the effort.


This exercise uses the deltoid, mainly the middle and posterior fibers, as well as the trapezius, triceps brachii, and serratus anterior. Although not worked as intensely, the rhomboids, infraspinatus, teres minor, and, deeper in, the supraspinatus also contract. You can also perform this exercise while standing at a frame that guides the barbell. Various specific machines can help with the performance of this exercise.

Attention: To prevent injury to the shoulder joint, which is vulnerable, lower the bar only as far as your unique shoulder structure and flexibility allow you to do comfortably (see page 39).



2

SEATED FRONT PRESSES



Sit with the back straight and hold the bar with an overhand grip, resting it across the upper chest:

- Inhale and extend the bar vertically.
- Exhale at the end of the movement.

This fundamental exercise mainly uses the anterior and lateral deltoids, clavicular head of the pectoralis major, triceps brachii, serratus anterior, trapezius, and, deeper in, the supraspinatus.


You can also perform this exercise standing, as long as you keep the back straight, avoiding excessive curvature of the lumbar spine.

Extending the barbell with the elbows forward isolates the anterior deltoid. Extending the bar with the elbows spread apart isolates the middle deltoid.

You can use various machines for this exercise.

SEATED DUMBBELL PRESSES

3




Sit on a bench, keeping the back straight, and hold dumbbells at shoulder level with an overhand grip (thumbs pointing inward):


- Inhale and extend the arms vertically.
- Exhale at the end of the movement.

This exercise contracts the deltoid, mainly the middle deltoid, as well as the trapezius, serratus anterior, and triceps brachii.

This movement may also be performed standing or alternating the arms. A backrest helps prevent an excessive arch in the back.



VARIATION
PALMS FACING EACH OTHER




FINAL POSITION



SHOULDER INJURIES




HIP JOINT



SHOULDER JOINT

Compared to the relatively stable coxofemoral joint, the shoulder joint is less encased and is more mobile, which makes it more vulnerable to injury.

ROTATOR CUFF MUSCLES FROM THE FRONT



Shoulder injuries occur frequently in weightlifting and especially in bodybuilding, where developing the entire deltoid group requires the athlete to perform a significant number of repetitions and variations in exercises, which multiplies the risk of injury.

Compared to the stability of the hip joint, where the head of the femur sits deep in the glenoid cavity of the pelvis, the shoulder joint, which is very mobile and allows the arm to move through a wide range of motion, is in fact much less contained and protected.

The shoulder is defined as a ball-and-socket joint because the head of the humerus is mainly held within the glenoid cavity of the scapula by a complex musculotendinous group.

Most weightlifting injuries occur when training the deltoids, and they rarely result in muscle pulls or tears. They are usually caused by poor technique or overuse of the tendons reinforcing the articular capsule.

In contrast to contact sports, such as football, where sudden arm movements can create serious injuries involving dislocation or even torn tendons, the most serious injury in weightlifting involves entrapment.

When some people perform exercises in which they raise the arms, such as extensions from the neck or lateral raises, the supraspinatus tendon is rubbed and compressed between the head of the humerus and the osteoligamentous ceiling created by the inferior surface of the acromion and the coracoacromial ligament.

Inflammation follows. This generally begins with the serous bursa, which normally protects the supraspinatus from excessive friction, and extends to the supraspinatus tendon itself, which, without treatment, ends up affecting the adjacent infraspinatus tendon posteriorly and the long head of the biceps brachii anteriorly. Raising the arm becomes extremely painful and


eventually can cause irreversible deterioration of the supraspinatus tendon through calcification and even tearing; however, this usually happens to people 40 years of age or older.

The space between the humerus and the osteoligamentous acromioclavicular ceiling varies from person to person. Some athletes cannot raise their arms laterally without excessive friction. These people should avoid all extensions from the neck, lateral raises that go too high, and back presses.

All barbell extensions for the shoulders must be performed to the front with the elbows slightly forward. When doing lateral dumbbell raises, you'll need to determine the proper height to raise the arms to. The correct movement is the one you can perform without causing pain.



Not everyone responds the same way to the same shoulder injury. Some people may perform all sorts of arm raises that compress the tendon, sometimes even causing tendon degeneration, without initiating a painful inflammatory process. This is how a torn supraspinatus tendon can be discovered during assessment without that person ever having complained of pain.

Another cause of shoulder pain may be an imbalance in muscle tension around the articular capsule. Remember that the head of the humerus is solidly fixed against the glenoid fossa of the scapula by a group of muscle tendons adhering to or crossing over the articular capsule: In front, this is the subscapularis; a little more anterior is the long head of the biceps; superiorly, is the supraspinatus; and finally posteriorly, the infraspinatus and teres minor. Spasm, hypertonicity, or hypotonicity in one or more of these muscles can pull the shoulder joint into an incorrect position. This position can cause friction during arm movements, resulting in inflammation.



Example: Shortening or spasm of the teres minor and the infraspinatus will pull the head of the humerus in external rotation, which will cause rubbing at the anterior shoulder joint during arm movement. Over time, this will injure the long head of the biceps brachii.


Balance the training of the shoulder muscles and avoid exercises that feel awkward or painful.



Comment: Massage, either manually or even better with an electric massager, and electrical stimulation are effective for decreasing or eliminating spasms and shortening of the teres minor and infraspinatus.



LYING DUMBBELL PRESSES



This is one of the rare exercises that may be performed by people suffering from the all-too-common entrapment syndrome.

Performing arm extensions with dumbbells while lying on a bench and keeping the elbows next to the body works the anterior deltoid and, to a lesser degree, the middle deltoid intensely while preventing excessive rubbing at the anterior shoulder.

When performed regularly, this maintains the size and tone of the deltoids despite the existence of injury. You can also use this exercise to reeducate the pectoralis major following tearing. Extending while keeping the elbows against the body reduces its stretch, thus reducing the risk of tearing the scarred area.

Performing the Exercise


Lie on a bench with the chest expanded, back slightly arched, feet flat on the ground, and the elbows bent next to the body, holding a dumbbell in each hand.

- Inhale and extend the arms vertically.
- Exhale at the end of the movement.
- Return to the initial position with a controlled movement.

HOW THE OSSEOUS MORPHOLOGY AFFECTS THE BACK PRESS




INFLUENCE OF THE LENGTH OF THE ARM ON LOWERING THE ELBOWS DURING THE BACK PRESS



- 1 Deltoid is stretched optimally, allowing for recruitment of the maximum number of muscle fibers during the initiation of the movement.
- 2 Deltoid is overstretched, which does not allow enough muscle fibers to be recruited for a powerful initiation of the movement.

INFLUENCE OF THE LENGTH OF THE CLAVICLES ON LOWERING THE BAR FROM THE BACK PRESS



It is important to take into account individual morphological differences when training the shoulders with the back press.

The length of the arms: The length of the arms, especially the forearms, plays a fundamental role in the execution of this exercise.

When lowering the bar below the ears, people with proportionately longer forearms bring their elbows down a lot lower than people with proportionately shorter forearms. This lower placement of the elbows strongly stretches the deltoid, putting it in an unfavorable position by not allowing the muscle to recruit the maximum number of fibers to initiate a powerful push.

To optimize training, people with proportionately longer arms should add heavier weights and not lower the bar too much below the ears. The most important issue is the intensity of feeling on the deltoids.

The length of the clavicles: The length of the clavicles has a considerable influence on the ability to lower the bar correctly behind the neck.

Short clavicles inevitably bring the shoulder blades toward the vertebral axis. During the extension from the neck, as both shoulder blades tilt to the center of the back (in adduction or external pendulum), the range of the shoulder blades is significantly reduced by the compression of the trapezius and rhomboid muscles in the middle of the back. This shortened range of the shoulder blades limits the ability to correctly lower the elbows in order to feel the deltoids working.

Furthermore, the greater the development of the muscles at the center of the back, the less the shoulder blades will be able to approach each other, which will further limit the lowering of the bar to the neck.

Comment: People with long forearms and short clavicles should avoid the back press in favor of better working the deltoids.





Attention

- 1 When the humerus rotates externally with elevation of the arm, reduced space between the glenohumeral joint and the osteoligamentous coracoacromial ceiling may lead to excessive friction that, over time, can damage or even tear the supraspinatus tendon. Therefore, at the slightest sensation of unease accompanied by pain during the back press, change the movement in order to avoid developing degenerative tendinitis of the rotator cuff.

- 2 Raising the arm (as in lateral raises with dumbbells) causes the humerus to internally rotate. Too narrow a space between the glenohumeral joint and the osteoligamentous coracoacromial ceiling may lead to excessive friction that risks damaging the infraspinatus tendon.

ROTATOR CUFF INJURIES WITH RAISING OF THE ARM




PINCHING OF THE SUPRASPINATUS WITH BACK PRESSES

During the back press, the supraspinatus tendon can get pinched between the acromion and the humerus.

4

ARNOLD PRESSES




Sit on a bench, keeping the back straight. With elbows bent and pointing forward, hold the dumbbells at shoulder level with an underhand grip (thumbs pointing away from each other):

- Inhale and extend the arms vertically while rotating 180 degrees at the wrists, bringing them into an overhand grip (thumbs pointing toward each other).
- Exhale at the end of the movement.

This exercise solicits the deltoid, mainly the anterior deltoid, as well as the clavicular head of the pectoralis major, triceps brachii, trapezius, and serratus anterior.

Variations: This exercise may be performed seated against a backrest to help prevent an excessive arch in the back, standing, and alternating arms.





Comment: Working with the elbows pointing forward prevents excessive friction, which triggers inflammation in the shoulder that can eventually develop into a more serious injury.


This movement is recommended for people with weak shoulders and is meant to replace more intense exercises, such as classic dumbbell extensions with the elbows pointing to the sides or extensions from behind the neck.

BENT-OVER LATERAL RAISES

5



DELTOID INSERTIONS




Stand with the legs slightly apart and knees slightly bent and lean forward at the waist while keeping the back straight. With arms hanging down, grasp the dumbbells with the elbows slightly bent:

- Inhale and raise the arms to horizontal.
- Exhale at the end of the effort.

This exercise works the shoulder group, accenting the work of the posterior deltoid. Squeeze the shoulder blades together at the end of the movement to contract the middle and lower portions of the trapezius, rhomboids, teres minor, and infraspinatus.


Variation: The exercise may be performed facedown on an incline bench.




FINAL POSITION

6

LATERAL DUMBBELL RAISES



SUPRASPINATUS MUSCLE ACTION




The supraspinatus helps the deltoid raise the arm laterally and helps keep the head of the humerus in the glenoid cavity.


Stand with a straight back, with legs slightly apart, arms hanging next to the body, holding a barbell in each hand:

- Raise the arms to horizontal with the elbows slightly bent.
- Return to the initial position.

This exercise mainly uses the middle deltoid. The three divisions of the deltoids create a multipennate muscle whose different fiber directions converge on the humerus. Their function is to support relatively heavy weight and to move the arm through its full range of motion with precision. Therefore, it is important to adapt training to the specifics of this muscle by varying the initial position of the movement (hands behind, to the side, or in front). This thoroughly works all the fibers of the middle deltoid. Because everyone's physical structure is different (length of the clavicle, shape of the acromion, and height of the insertion at the humerus), you must find the angle of the initial position that is best for you. Lateral raises contract the supraspinatus, although you can't see this because it is located deep in the supraspinatus fossa of the scapula (shoulder blade), where it attaches to the lesser tubercle of the humerus.


Raising the arm above horizontal contracts the upper part of the trapezius; however, many bodybuilders don't work above horizontal so that they isolate the lateral deltoid. This exercise should not be performed with heavy weights, but instead in sets of 10 to 25 reps, while varying the working angle without much recuperation time until you feel a burn. To increase the intensity, maintain an isometric contraction for a few seconds with the arm at horizontal between each repetition.





**CONVERGENT
MUSCLE**

**PENNATE
MUSCLE**




**CONVERGENT
MUSCLE**

**PENNATE
MUSCLE**

A pennate muscle displaces greater weight than a convergent muscle but over a shorter distance. During lateral raises, because the pennate fibers of the middle deltoid produce great force but weak contraction, they work synergistically with the anterior and posterior deltoids to bring the arm to horizontal.

The sum of the actin* and myosin* filaments of a convergent muscle is equal to its transverse section (A).
The sum of the actin and myosin filaments of a pennate muscle is equal to the sum (A) of its oblique sections (A1 and A2).

*The motor elements of a muscle whose maximum force of contraction is approximately 5 kg per cm² per second.





FINAL POSITION: VARIATIONS

- 1 Arms raised to horizontal works the deltoid.
- 2 Arms raised above horizontal isolates the upper and middle portions of the trapezius.

7

ALTERNATE FRONT ARM RAISES



Stand with the feet slightly apart. Hold the barbells with an overhand grip as they rest against the front of the thighs or slightly to the side:

- Inhale and alternate raising the arms to the front to eye level.
- Exhale at the end of the effort.

This exercise uses mainly the anterior deltoid, the clavicular head of the pectoralis major, and, to a lesser degree, the remaining deltoids. All movements that raise the arms contract the muscles that anchor the scapula to the rib cage, such as the serratus anterior and rhomboids, which create a stable support for the humerus to move from.




VARIATION
LYING FACEDOWN ON AN INCLINE BENCH



VARIATION
RAISING TO THE FRONT
USING BOTH HANDS

SIDE-LYING LATERAL RAISES


8



Lie on one side on the floor or on a bench holding a dumbbell with an overhand grip:

- Inhale and raise the arm to vertical.
- Exhale at the end of the movement.


Unlike standing raises, which progressively work the muscle to maximum intensity at the end of the movement (when the arm reaches horizontal), this exercise works the deltoid differently by focusing the effort at the beginning of the raise. Sets of 10 to 12 repetitions work best.



Comment: This movement contracts the supraspinatus, the muscle mainly responsible for initiating abduction. Varying the initial position (dumbbell in front of or behind the thigh) allows you to work all the deltoid fibers. To increase the intensity of the movement, perform this exercise with continuous tension without resting the dumbbell on the thigh.

9


LOW-PULLEY FRONT RAISES, OVERHAND GRIP



Stand with the feet slightly apart, arms next to the body. Grasp the handle with an overhand grip with one hand:


- Inhale and raise the arm up to eye level.
- Exhale at the end of the movement.

This exercise contracts the deltoid (mainly the anterior deltoid) as well as the clavicular head of the pectoralis major and, to a lesser degree, the short head of the biceps brachii.





LOW-PULLEY FRONT RAISES, NEUTRAL GRIP

10



FINAL POSITION

THE MOVEMENT
WITH A DUMBBELL

Stand with legs slightly apart and your arm by your side holding the handle with a neutral (semipronated) grip (this exercise is performed with a handle adapted for the neutral grip):

- Inhale and raise your arm forward up to eye level; exhale at the end of the raise.
- Slowly return to the initial position and repeat.

This exercise mainly works the anterior deltoid as well as the clavicular part of the pectoralis major and, to a lesser degree, the middle deltoid and the short head of the biceps.


It is preferable to work this movement in a long series.

Variation: Perform the exercise with a harness.

Comment: This exercise is excellent for people who have difficulty developing the anterior deltoid. The semipronated grip engages the humerus in external rotation, which at the beginning of the movement stretches the anterior fibers of the deltoid, allowing you to feel them working.





STRETCHING THE ANTERIOR DELTOIDS



Stand with arms shoulder-width apart and extended behind your back with one hand grasping the other. Push the arms as far back as possible, then slowly raise them while pushing out the chest and pulling the chin in. Hold this position for 10 seconds. This exercise mainly stretches the anterior deltoid as well as the pectoralis major and biceps brachii. The brachialis, brachioradialis, and extensor muscle group at the wrist are also recruited.

HIGH-PULLEY LATERAL EXTENSIONS

11



RHOMBOID MAJOR AND MINOR MUSCLES


Located deep under the trapezius, the rhomboids pull the shoulder blades together toward the spine and press them against the rib cage.

In some people, the major and minor rhomboids are fused, creating one muscle.

Stand facing the pulleys with the arms extended to the front. Grip the right handle with the left hand and the left handle with the right hand:

- Inhale and extend the arms to the side and back.
- Exhale at the end of the movement.
- Return to the initial position with a controlled movement and begin again.

This exercise mainly contracts the posterior deltoid, infraspinatus, teres minor, and, at the end of the movement as the shoulder blades come together, the trapezius and, deeper in, the rhomboids.







Comment: People who carry their shoulders forward because of chest muscle development can perform this exercise in addition to posterior shoulder work at a machine to help rebalance their posture.

To realign the shoulders where they belong, work with moderate weights and squeeze the shoulders back at the end of the movement.



12 EXTERNAL ARM ROTATIONS AT A PULLEY



Position the pulley at waist level and position your body in line with the apparatus. Grip the handle with your forearm in front of your body, your elbow bent, and your upper arm against your body:

- Externally rotate your arm, trying to keep the upper arm against your body with the elbow bent.

This exercise mainly works the infraspinatus and teres minor as well as the posterior deltoid. If at the end of the movement the scapula is brought to the center of the body, the rhomboids and the middle and inferior portions of the trapezius will be worked at the same time.



This movement is mainly used to strengthen the infraspinatus and prevent painful contractures and frequent injuries of this muscle.

External rotations of the arm at the pulley are often recommended during recovery from a tear or partial tear of the infraspinatus. Use very light weights initially.

Comment: Perform this exercise with the objective of working the posterior deltoid, which is often difficult to recruit. In this case, move your arm slightly away from the body and extend the elbow at the end of the movement.

LOW-PULLEY BENT-OVER LATERAL RAISES

13



PARTS WORKED


Stand with the feet apart and legs slightly bent. Lean forward from the waist, keeping a flat back. Grip a handle in each hand with the cables crossed:

- Inhale and raise the arms to the sides to horizontal.
- Exhale at the end of the effort.

This exercise mainly works the posterior deltoid. At the end of the movement, as the shoulder blades squeeze together, the trapezius (middle and lower portions) and the rhomboids contract.



14 LOW-PULLEY LATERAL RAISES





Sternocleidomastoid
Scalenes
Trapezius
Omohyoid

Deltoid
– Anterior deltoid
– Middle deltoid
– Posterior deltoid

Biceps brachii
Brachialis
– Long head
– Lateral head
– Medial head


Brachioradialis
Extensor carpi radialis longus
Anconeus
Extensor carpi ulnaris
Flexor carpi ulnaris
Extensor carpi brevis
Extensor digitorum

Extensor digiti minimi
Abductor pollicis longus
Extensor pollicis brevis



Pectoralis major
Rectus abdominis, under the aponeurosis
External oblique

DELTOID ACTION



Grasp the handle with the arm next to the body:

- Inhale and raise the arm to horizontal.
- Exhale at the end of the movement.

This exercise mainly develops the middle deltoid. Because the muscle is multipennate, composed of many fibers in the shape of a feather, it is best to vary the working angles in order to work all the fibers.




FINAL POSITION

IMPORTANCE OF PULLING THE SHOULDERS BACK



MUSCLES ENGAGED WHILE PULLING THE SHOULDERS BACK



One of the major postural defects encountered most frequently in modern society, where we spend inordinate amounts of time in the sitting posture, is kyphosis (rounding of the upper thoracic spine).

This bad position of the upper body is most often due to the hypotonic state of the muscles approximating the shoulder blades and the external rotators of the arms, or, more frequently with men, the hypertonicity and predominance of the development of the chest muscles. In weight training, focusing extensively on the pectorals or overtraining in bench presses can also contribute to this postural defect.


It is important to perform postural rebalancing by practicing specific exercises to straighten the shoulders, such as the pec deck rear-delt laterals (page 58), external arm rotations at a pulley (page 50), or bent-over lateral raises (page 41).



Rounding the upper back during deadlifts limits the power of the lift. To avoid this rounding of the back, perform specific exercises to strengthen the muscles responsible for straightening the shoulders.

Comment: With powerlifting and heavy deadlifts, it is essential to avoid rounding of the shoulders to the front and excessive cantilevering, which can restrict the power of the lift. Always straighten the shoulders during the execution of the movement and use specific exercises to prepare for powerlifts and heavy deadlifts.

15 ONE-DUMBBELL FRONT RAISES




Stand with the legs slightly apart, back straight, and the abdominal muscles contracted. With arms extended, grasp a dumbbell in both hands with fingers crossed over each other and rest it against the thighs:

- Inhale and raise the dumbbell to eye level.
- Lower gently, avoiding abrupt movements.
- Exhale at the end of the movement.

This exercise mainly contracts the anterior deltoid, the clavicular head of the pectoralis major, and the short head of the biceps.


Note that all the fixators of the scapula are used during the isometric contraction, which allows the humerus to move from a stable position.



Clavicular head of the pectoralis major actively participates in raising the arm anteriorly.

BARBELL FRONT RAISES

16



VARIATION
FRONT RAISE AT A LOW PULLEY

Stand with the legs slightly apart and the back straight, contracting the abdominal muscles. Hold the barbell with an overhand grip and rest it against the thighs:

- Inhale and raise the barbell with extended arms to eye level.
- Exhale at the end of the movement.


This exercise contracts the anterior deltoid, the clavicular head of the pectoralis major, the infraspinatus, and, to a lesser degree, the trapezius, serratus anterior, and short head of the biceps.

If you continue raising the arms, the posterior deltoid contracts, reinforcing the work of the other muscles and allowing you to raise the arms to vertical.

The exercise may also be performed with your back to a low pulley and the cable passing between the legs.

1 Begin

2 End




THE MOVEMENT

Comment: The biceps brachii participates to a lesser degree in all anterior arm raises.

17

UPRIGHT ROWS




Stand with the legs slightly apart and back straight. Grasp the barbell with an overhand grip slightly wider than shoulder width and rest it against the thighs:

- Inhale and pull the barbell up along the body to the chin, keeping the elbows as high as possible.
- Lower the bar in a controlled manner without abrupt movements.
- Exhale at the end of the effort.


This exercise mainly uses the deltoid, trapezius, and biceps, and, to a lesser degree, the muscles of the forearms, the gluteal muscles, the lumbosacralis group, and the abdominal muscles.

This is a fundamental exercise that is comprehensive and helps develop a "Hercules" physique.



MACHINE LATERAL RAISES

18




Sit at a machine and grasp the handles:




- Inhale and raise the elbows to horizontal.
- Exhale at the end of the movement.

This exercise uses the deltoid (focusing most of the effort on the middle deltoid) and the supraspinatus, located under the deltoid. If raising the arm above horizontal, the upper portion of the trapezius also becomes involved.

Comment: This is an excellent beginning exercise because you don't have to worry about your form or positioning, and it allows long sets of repetitions.




19 PEC DECK REAR-DELT LATERALS



INFRASPINATUS AND TERES MINOR

Both of these muscles arise from the posterior surface of the scapula (shoulder blade), pass onto the scapulohumeral articulation (adhering to its articular capsule), and insert at the greater tubercle of the humerus. They play an important role in external rotation of the arm and reinforce the action of the shoulder ligaments by actively reinforcing the attachment of the arm to the chest.

Comment: In some people, the teres minor and infraspinatus are fused, forming one muscle.




THE MOVEMENT

Sit facing the machine, with the torso against the back pad and arms forward, gripping the handles:

- Inhale and separate the arms, squeezing the shoulder blades together at the end of the movement.
- Exhale.

This exercise mainly engages the posterior deltoid, infraspinatus, and teres minor and, at the end of the movement when the shoulder blades squeeze together, the trapezius and rhomboids.

STRETCHING THE POSTERIOR ROTATOR CUFF MUSCLES




Stand with a dumbbell in your hand, with your torso cantilevered forward and resting on the machine (for example, the arm rest of a Larry Scott desk or on a bench with lumbar extensions).

With your arm released, let the dumbbell hang for a minute while trying to relax the shoulder.

This exercise allows you to stretch the infraspinatus, teres minor, and, to a lesser extent, the supraspinatus; these muscles attach to the posterior surface of the scapula. In weightlifting, these muscles are often the site of contractures and spasms that engage the shoulder in a bad position, which over time may lead to particularly incapacitating tendon overuse pathologies.

Comment: Contractures or spasms of the teres minor and infraspinatus will engage the humerus in external rotation. This creates excessive friction on the long head of the biceps at the front of the arm (in the bicipital groove). If not treated, this can lead to inflammation and tearing of the tendon. Therefore, at the slightest suspicion of a contracture, it is important to relax these muscles while performing this specific stretch.





THE MOVEMENT



STRETCHING THE SHOULDER

THE TWO WAYS OF PULLING ON THE ELBOW



Stand with your head level and one arm horizontal. Grasp the elbow with the opposite hand and pull on the arm to slowly bring the elbow toward the opposite shoulder.

Maintain this position for 10 to 20 seconds, the time it takes to properly feel the stretch.

This exercise mainly works the posterior fascicle of the middle deltoid and the teres minor and infraspinatus. These small external rotator muscles of the humerus are frequently the site of contractures, which can lead to functional disequilibrium of the shoulder (such as excessive friction of the tendon of the long head of the biceps in the bicipital groove of the humerus) and may result in pathological inflammation.

The middle and inferior portions of the trapezius muscle and the rhomboid major are also stretched.

Variation: Pull the elbow with the opposite arm passing underneath.

Comment: For some people with very well-developed muscles, adduction of the arm can be hindered by compression of the biceps brachii against the pectoralis major, which will limit the stretch at the posterior part of the shoulder.



3

CHEST


| | |
|---|-----------|
| 1 INCLINE BENCH PRESSES | 62 |
|  STRETCHING THE PECTORALIS MAJOR | 63 |
| 2 BENCH PRESSES | 64 |
|  ACROMIOCLAVICULAR PROBLEMS | 66 |
|  PECTORALIS MAJOR TEAR | 68 |
| 3 CLOSE-GRIP BENCH PRESSES | 69 |
|  BENCH PRESSES AND ELBOW PAIN | 69 |
|  BENCH PRESSES AND MORPHOLOGY | 70 |
| 4 DECLINE BENCH PRESSES | 72 |
| 5 MACHINE BENCH PRESSES | 73 |
| 6 PARALLEL BAR DIPS | 74 |
|  PAY ATTENTION TO THE NECK POSITION! | 75 |
| 7 PUSH-UPS | 76 |
| 8 DUMBBELL PRESSES | 77 |
| 9 DUMBBELL FLYS | 78 |
| 10 INCLINE DUMBBELL PRESSES | 79 |
| 11 INCLINE DUMBBELL FLYS | 80 |
| 12 PEC DECK FLYS | 81 |
| 13 CABLE CROSSOVER FLYS | 82 |
| 14 DUMBBELL PULLOVERS | 83 |
| 15 BARBELL PULLOVERS | 84 |

1

INCLINE BENCH PRESSES



Comment: Contrary to popular lore, the incline press does not tone the breasts and in no way prevents their sagging. Breasts are composed of adipose tissue containing the mammary glands, all of which is contained in a net of connective tissue that rests on top of the pectoralis major.




Sit on an incline bench angled at 45 to 60 degrees and grasp the barbell with an overhand grip wider than shoulder width:

- Inhale and lower the barbell to the sternal notch.
- Extend the arms.
- Exhale at the end of the movement.

This exercise mainly solicits the clavicular head of the pectoralis major, anterior deltoid, triceps brachii, serratus anterior, and pectoralis minor. This exercise may be done at a frame that guides the bar.

STRETCHING THE PECTORALIS MAJOR



Stand with your arm extended. Grip a support with your hand and slowly rotate your chest to the opposite side, away from the support.


This exercise mainly stretches the pectoralis major, anterior deltoid, and biceps brachii.

Variation: Position your hand at various levels in order to stretch all the fascicles of pectoralis major.

Comment: This is an excellent stretch for the bench press in weightlifting and all throwing sports, including tennis, volleyball, and handball.

2


BENCH PRESSES



Lie faceup on a horizontal bench, with buttocks on the bench and feet flat on the ground:

- Grasp the barbell with an overhand grip wider than shoulder width.
- Inhale and lower the bar to the chest with a controlled movement.
- Extend the arms and exhale at the end of the effort.


This exercise engages the complete pectoralis major muscle, pectoralis minor, anterior deltoid, serratus anterior, and coracobrachialis.




THE MOVEMENT

Variations

1. This movement may be performed while arching the back powerlifter style. This position brings the more powerful lower part of the pectoral muscle into play, allowing you to lift heavier weights.
2. Executing the extension with the elbows next to the body concentrates the work onto the anterior deltoid.
3. Varying the width of the hands isolates different parts of the muscle:
 - Hands closer together isolates the central part of the pectorals.
 - Hands wider apart isolates the lateral part of the pectorals.
4. Varying the angle of the barbell isolates different parts of the muscle:
 - Lowering the bar to the chondrocostal border of the rib cage isolates the lower part of the pectorals.
 - Lowering the barbell onto the middle part of the pectorals isolates the midline fibers.
 - Lowering the bar onto the sternal notch isolates the clavicular head of the muscle.
5. If you have back problems or want to isolate the pectorals, perform the extension with the legs raised.
6. Perform the extension at a frame that guides the bar.




CLASSIC POSITION



ARCHED-BACK VARIATION


Executing the bench press with an arched back, powerlifter style, limits the range of the movement and allows you to lift significantly heavier weights because it uses mainly the lower part of the pectorals, which are the strongest. In competition, the feet and the head should not move, and the buttocks should remain in contact with the bench. People with back problems should not perform this variation.




RAISED-LEG VARIATION

Performing the movement with raised legs helps prevent excessive arching, which can cause low back pain.

This variation diminishes the effort of the lower pectorals by working the middle and superior fibers more.



POSITIONING FOR A POWER BENCH PRESS




ATTENTION

- 1 For maximum safety, lock onto the bar with a grip in which the thumb and fingers oppose each other.
- 2 If the grip on the bar is not locked on in opposition, the bar could slip out of your hands and fall on the jaw, or worse, the neck, and cause a serious injury.



ACROMIOCLAVICULAR PROBLEMS




Acromioclavicular injuries are common in weightlifting. Most dedicated weightlifters encounter this problem.

Unlike other sports (such as rugby, football, and riding) and combat sports with throws (wrestling and judo) where the shoulder joint can become severely injured with violent contact or a fall involving acromioclavicular dislocation with ligament tearing, in weightlifting, acromioclavicular pathologies are mainly due to microtraumas resulting from excessive and repetitive stress and poor control of the shoulder articulation.

Pain develops progressively. Although pain is easily tolerated at the beginning, it gradually disturbs the practice of weightlifting until it finally hinders the ability to perform a great number of exercises such as bench presses and dips. All downward propulsions become painful, and leaning on the elbows might also be painful. Examination of the acromioclavicular articulation reveals slight swelling and pain on palpation. Although not really serious, this type of injury generally takes a long time to heal. It also takes a long time for the inflammation to subside and the articular capsule with the stretched acromioclavicular ligaments to revert to their normal size and allow for normal joint play.

When this injury occurs, training of the upper body must stop for two weeks.


CROSS SECTION OF THE ACROMIOCLAVICULAR JOINT



*The acromioclavicular ligament is a thickening of the superior part of the articular capsule of the acromioclavicular joint.

Comment: At the osseous level, the upper extremity is attached to the chest by the clavicle, which extends from the sternum to the shoulder blade. Although not very mobile, the clavicular articulations are often overused and subject to inflammatory wear pathologies.

BASIC ANATOMY OF THE CHEST




On resuming training of the upper body, avoid for at least two months the bench press and all movements that involve pushing down, such as decline presses and dips, since they risk stretching the acromioclavicular ligaments all over again. On the other hand, all movements that involve pushing up, such as incline presses or presses for the shoulders with barbells and dumbbells, can be performed without risk because they tend to stabilize the acromioclavicular articulation, which limits the risk of stretching the ligaments.

Disregarding this advice will cause the articular inflammation to continue and, in certain people, will lead to intra-articular calcifications and may seriously compromise an athletic career.


Prevention: In weightlifting, acromioclavicular inflammations most often appear after overtraining of the bench press with series that are too long or poorly controlled (rapid lowering, jerking, and bouncing on the chest). Powerlifters who perform the bench press are also susceptible to tension generated on the acromioclavicular ligament that initiates inflammatory pathologies.

As soon as you feel any pain, stop these traumatizing techniques for some time and replace them with exercises for the pectorals, such as standing spreads at the pulley in conjunction with movements using dumbbells. Always work with a certain degree of incline on the bench.





PECTORALIS MAJOR TEAR



The pectoralis major originates at the anterior surface of the rib cage and inserts at the anterior surface of the upper end of the humerus.

It is a powerful muscle whose main function is to bring the arms together in front of the rib cage. (It is the hugging muscle.)

Unlike most sports, where pectoralis major injuries are rare, weightlifting, especially the bench press, can lead to small tears and even partial rupture of its tendon.

This ultimate injury is seen only in relatively powerful athletes using abnormally rapid force before the tendon

has had time to strengthen. Sometimes it is associated with a low-calorie diet aimed at increased muscle definition. (These diets tend to weaken the muscles, tendons, and joints.)

The injury, which always occurs during heavy bench-pressing, generally affects only the tendon of the clavicular head of the pectoralis major.


A torn tendon is extremely painful, and the athlete may faint. Swelling and bruising often appear on the anterior surface of the arm, and retraction of the clavicular head leads to a hollow that is medial to the anterior deltoid.

The problem with this injury is that doctors often misdiagnose it. This mistake is unfortunately common but is understandable because during the post-traumatic examination the injured party is able to perform all the movements that indicate full motor function of the pectoralis major. Therefore, the injury appears to be a simple muscle tear rather than the more serious tear of the tendinous insertion.

For example, despite a tear of the clavicular head of the pectoralis major, anterior elevation of the arm, which is part of its function, is compensated for by the anterior deltoid. And abduction is performed by the sternal and abdominal heads of the pectoralis major.

If the tendon of the clavicular head of the pectoralis major is torn, it must be surgically reinserted onto the humerus as soon as possible. If this is not done promptly, retraction and fibrosis of the muscle occurs, and the operation will no longer be possible.

Although you can move your arm through its full range of motion without the superior head of the pectoralis major, you will never recover your initial strength and will be at a serious disadvantage if you want to continue heavy weight training.



INSERTION OF THE PECTORALIS MAJOR MUSCLE ON THE HUMERUS DISPLAYING HOW THE TENDON TWISTS ON ITSELF CREATING A U-SHAPE


During bench presses or flies, the most lateral part of the pectoralis major tendon, which corresponds to the clavicular head, is put under the most stress.

Therefore, when lifting heavy weights, this is the tendon that tears or pulls away from its insertion.

Insertion of the pectoralis major

CLOSE-GRIP BENCH PRESSES

3



Attention: Depending on your physical structure, the narrow grip may cause wrist pain. In this case, use a wider grip.

Lie on a horizontal bench with the buttocks on the bench and the feet on the ground. Grip the barbell with an overhand grip and wrists 4 to 15 inches apart, depending on the flexibility of the wrists:

- Inhale and lower the bar with a controlled movement to the chest, with the elbows out to the sides.
- Extend and exhale at the end of the effort.

This exercise develops the pectoral muscles at the sternal notch and the triceps brachii. (With this in mind, it may be included in a program for the arms.) By extending and keeping the elbows next to the body, a greater part of the work is performed by the anterior deltoid. This movement may be performed at a frame that guides the bar.



Bench Presses and Elbow Pain


Elbow pain most often develops after bench pressing. This overuse injury is generally related to excessive training with long sets. In bench pressing, locking the extended arms at the end of the movement subjects the elbow to rubbing and microtrauma, which over time may lead to inflammation.

Comment: Occasionally, this condition can lead to intra-articular calcifications, which are particularly crippling. In this case, surgery is often the only solution for regaining complete arm extension.

At the first sign of elbow pain, avoid for several days exercises that involve arm extension in order to prevent serious injury.

When you resume exercises that include arm extension, avoid completely extending the forearms at the end of the movement until the pain has completely disappeared.

SECTION THROUGH THE ELBOW JOINT



With repeated extension of the forearm, the olecranon butts up against the olecranon fossa of the humerus. The articulation then suffers from microtrauma, which over time may generate painful inflammation at the dorsal surface of the elbow.



BENCH PRESSES AND MORPHOLOGY

The bench press is by far the most-used exercise in weightlifting. It is also the exercise that causes the most injuries. Therefore, to perform this movement correctly and to reduce risk, you must learn the basics of individual morphological differences.

Arm Length

Besides wear-and-tear pathologies, most injuries are muscle tears or tendon tears of the pectoralis major. These occur most often during the negative phase of the movement—that is, while lowering the barbell. In lowering the barbell to the chest, the pectoralis major, which inserts onto the humerus, becomes increasingly stretched and vulnerable as the arm is lowered.




But the lowering of the arm and the stretch of the pectoralis major vary significantly depending on the individual. The longer the arm and especially the forearm, the lower the humerus will go down, and the more the pectoralis major will be dangerously stretched. Therefore, it is not surprising that most injuries occur in people with relatively longer arms.

Thickness of Rib Cage

The thicker the rib cage, the more the lowering of the barbell and by extension the stretching of the pectoralis major will be restricted.

People with a thick rib cage will theoretically perform the bench press without much risk of injury to the pectoralis major. It is therefore not surprising that most of the great bench press champions are shorter-limbed people (that is, people with proportionately shorter extremities and barrel chests). These two details allow them to achieve their records relatively safely.

Remember that injury is what often limits athletes' progress. In addition to training methods, nutrition, and mental state, individual morphology plays a fundamental role in success in sports. It is therefore essential to adapt your training to your morphology and to understand that what is good for one person may not be as good for another.




Limiting the Risk of Injury

It is possible to perform the bench press by bringing the hands closer together. Because this variation limits the lowering of the arms, it reduces the stretch of the pectoralis major and limits the risk of injury.


Although the amplitude of the movement is greater, the work on the biceps is more intense, and the performance is reduced, this low-risk variation is sometimes used by certain long-limbed champions of the bench press.

To avoid excessive stretching of the pectoralis muscles, it is possible to partially perform the bench press by shortening the lowering phase of the bar so that it does not touch the chest.



1 Performing the bench press with elbows apart

2 Performing the bench press with elbows closer together



Muscle Prominence

Note that there are two ways of performing the bench press according to the strong features of individual muscles.

The bench press can be performed with the elbows apart. This technique directs most of the effort onto the pectoralis major.

The bench press can also be performed with the elbows closer together, closing the angle of the arm and chest. People who have deltoids that are stronger than the pectoralis major will instinctively use this technique.


Aside from the morphology, both of these types of bench presses can be used to direct the work onto the pectoralis major (elbows spread) or onto the deltoids (elbows closer together).



Attention: In the supine bench press, it is important to take the individual morphology into account.


A: A small rib cage combined with long arms increases the excursion of the bar, making the movement painful while at the same time limiting the strength developed. Furthermore, when the bar approaches the chest, the pectoralis major is dangerously stretched. When heavier weights are used, it increases the risk of muscular tears and tendinous disruptions.

B: A barrel chest coupled with short arms allows for safe performance of a supine bench press by limiting the amplitude of the movement and the stretch of the pectoralis major at the end of the descent of the bar (when the bar touches the chest). It is not surprising that the greatest champions of the supine bench press have this type of morphology.



4

DECLINE BENCH PRESSES




Lie on a decline bench (between 20 and 40 degrees), with the head angled down and feet fixed to prevent sliding. Grasp the barbell with an overhand grip shoulder width or more:




- Inhale and lower the barbell to the lower pectorals with a controlled movement.
- Extend the arms and exhale at the end of the movement.

This exercise contracts the pectoralis major (mainly its inferior fibers), triceps brachii, and the anterior deltoid.

This exercise is useful for outlining the inferior groove of the pectorals. Using light weights and lowering the bar to the neck stretches the pectoralis major correctly. The decline press may be performed at a frame that guides the bar.




PECTORALIS MAJOR

PECTORALIS MAJOR
MUSCLE INSERTIONS

PART OF THE PECTORAL
MUSCLES MAINLY USED

MACHINE BENCH PRESSES

5




Lie on the machine with the gluteals in contact with the bench and feet flat on the floor. Grip the handles:

- Inhale and press.
- Exhale at the end of the movement.

This guided exercise, reminiscent of the dumbbell press (page 77), focuses the effort on the pectoralis major, mainly on the sternal parts at the end of the execution.


The triceps and the posterior fascicles of the deltoids are also recruited, although less intensely.

Variation: When arching the back and pushing the chest out, part of the effort is put onto the inferior fascicles of the pectoralis major, but this technique is contraindicated for people who have back pain.



6

PARALLEL BAR DIPS



Hang from the parallel bars with arms extended and legs suspended:


- Inhale and bend the elbows to bring the chest level with the bars.
- Return to the extended arm position.
- Exhale at the end of the effort.

The more the chest is angled forward during the exercise, the more the inferior fibers of the pectorals are used. Conversely, the more vertical the chest, the more the triceps brachii will be used. This exercise is excellent for stretching the pectoralis major and for working on the flexibility of the shoulder girdle. However, it is not recommended for beginners because it requires a certain amount of initial strength.

If you are a beginner, use a dips machine to familiarize yourself with the movement.


Sets of 10 to 20 repetitions provide the best results.

For developing more strength and also more size, athletes used to this movement may use a weight belt, or hang a weight from their legs.



EXECUTING DIPS AT A MACHINE

- 1 Initial position
- 2 Final position




THE MOVEMENT

Comment: Execute the dips with caution to prevent shoulder trauma.

PAY ATTENTION TO THE NECK POSITION!



DIAGRAM OF THE NERVES OF THE UPPER EXTREMITY



In weightlifting, a faulty position of the neck during certain exercises may lead to bothersome and incapacitating neuralgia in people predisposed to the condition.

These neuralgias manifest as numbness in the arm accompanied by the sensation of pins and needles and sometimes local insensitivity.

These symptoms most often appear in the days after the execution of dips (page 74), pec deck rear-delt laterals (page 58), squats (page 126), and deadlifts (page 104) when these movements are done with the neck in extension and the head thrown back.

In fact, throwing the head back can initiate spasms and contractions of the deep muscles of the neck, leading to compression of the spinal nerves as they exit from the cervical vertebrae.


This compression involves a neuralgia that most often affects the brachial plexus at the vertebral levels of C4, C5, C6, C7, C8, and T1 (C stands for cervical and T for thoracic).

To find out where the involved nerve exits the spine, you need only look at the diagram, then follow the nerve from its pins and needles and numbness and ascend to its vertebra.

To avoid these types of neuralgia, perform dips or reverse push-ups at a machine by bringing the head forward while at the same time bringing the chin to the chest.

With the squat or deadlift, perform the exercise while keeping the neck very straight and looking forward.



If neuralgia has manifested, stop performing the exercise with the head thrown backward and the neck in extension.



Attention: Executing dips with the neck in extension can cause neuralgia in some people.

7

PUSH-UPS



INITIAL POSITION

Support yourself facedown on the ground, with arms extended, hands shoulder-width (or more) apart, and feet touching or slightly apart:

- Inhale and bend the elbows to bring the rib cage close to the ground without arching the low back excessively.
- Push back up to complete arm extension.
- Exhale at the end of the movement.

This movement is excellent for the pectoralis major and the triceps brachii.


Variations

Varying the tilt of the chest focuses the work on different parts of the pectorals:


- Feet higher isolates the clavicular head of the pectoralis major.
- Chest higher isolates the inferior part of the pectoralis major.

Varying the width of the hands focuses the work on different parts of the pectorals:

- Hands wider isolates the lateral part of the pectoralis major.
- Hands closer together isolates the sternal head of the pectoralis major.






PART OF THE PECTORAL MUSCLES MAINLY USED



While performing push-ups the serratus anterior contracts to maintain the scapula against the rib cage, locking the arms onto the torso.

DUMBBELL PRESSES

8




Lie faceup on a horizontal bench, with feet flat on the ground for stability and elbows bent. Hold the dumbbells with an overhand grip at chest level:


- Inhale and extend the arms vertically while rotating the forearms so that the palms face each other.
- Once the hands face each other, perform an isometric contraction to focus the effort on the sternal head of the pectoralis major.
- Exhale at the end of the movement.

This exercise is similar to the bench press, but with its greater range of motion, it stretches the pectoralis muscles.

Although not contracted as intensely, the triceps brachii and anterior deltoid are also used.




VARIATION
EXECUTING THE EXERCISE
WITHOUT ROTATING THE FOREARMS



PART OF THE PECTORAL
MUSCLES MAINLY USED

9

DUMBBELL FLYS




Lie on a narrow bench that won't interfere with the shoulder movement and hold a dumbbell in each hand with arms extended or slightly bent to relieve stress on the joint:


- Inhale and open the arms to horizontal.
- Raise the arms to vertical while exhaling.
- Perform a small isometric contraction at the end of the movement to emphasize the work on the sternal head of the pectoralis major.

This exercise is never performed with heavy weights.

This exercise focuses the work on the pectoralis major. It serves as a basic exercise to increase thoracic expansion, which contributes to increased pulmonary capacity. It also develops muscle flexibility.



THE MOVEMENT




PART OF THE PECTORAL MUSCLES MAINLY USED



Attention: To avoid the risk of tearing the pectoral muscles, perform the exercise with extreme caution when using heavier weights.

INCLINE DUMBBELL PRESSES

10




Sit on a bench with an angle of no more than 60 degrees (to prevent too much work with the deltoid), with elbows bent. Grasp the dumbbells with an overhand grip:

- Inhale and extend the arms vertically, bringing the dumbbells together.
- Exhale at the end of the movement.


This exercise, which is midway between an incline press and incline dumbbell fly, works the pectorals (mainly the clavicular head) and increases their flexibility. It also contracts the anterior deltoid, the serratus anterior, and the pectoralis minor (these last two muscles are fixators of the scapula, which stabilize the arm at the torso). It also uses the triceps brachii, but not as intensely as the barbell press does.

Variation: Beginning the press with the hands in an overhand grip and rotating the wrists halfway through the movement so that the dumbbells face each other focuses the effort on the sternal head of the pectoralis major.



11


INCLINE DUMBBELL FLYS




Sit on a bench angled between 45 and 60 degrees, dumbbells in hand and arms extended vertically or slightly bent to ease stress when bringing the arms together:

- Inhale and extend the arms to horizontal.
- Raise the arms to vertical while exhaling.

This movement should not be performed with heavy weights. It focuses the effort mainly on the clavicular head of the pectoralis major. Along with the pullover, it is a fundamental exercise for developing thoracic expansion.




PART OF THE PECTORAL MUSCLES MAINLY USED



FINAL POSITION

PEC DECK FLYS

12






Sit at the machine with the arms open and horizontal, bent at the elbows. Rest the forearms on the pads, with the forearms and wrists relaxed:

- Inhale and squeeze the arms together.
- Exhale at the end of the movement.


This exercise works the pectoralis major by stretching it. As the elbows come together, focus the effort onto the sternal head of the pectoralis major.

This exercise also develops the coracobrachialis and the short head of the biceps brachii. Long sets allow you to pump the muscles intensely.



This exercise helps beginners develop enough strength to move onto more complex movements.



13 CABLE CROSSOVER FLYS




PART OF THE PECTORAL MUSCLES MAINLY USED



Stand with the legs slightly apart and lean the torso forward a bit, with the arms spread apart and elbows slightly bent:


- Inhale and squeeze the arms together until the wrists touch.
- Exhale at the end of the contraction.
- Return without jerking to the initial position and repeat.

This is an excellent exercise for working the pectoralis major muscles. Sets with a lot of reps allow you to pump the muscle well. You can work all the fibers of the pectoralis major by varying the angle of the chest and the working angle of the arms (squeezing the arms at various heights).




THE MOVEMENT

Comment: Cable crossover flys also contract the pectoralis minor, which is located deeper than the pectoralis major. Besides stabilizing the scapula (shoulder blade), this muscle also pulls it forward.




- 1 Crossed arms
Increases the focus onto the sternal head of the pectoralis major
- 2 Classic execution



INITIAL POSITION

DUMBBELL PULLOVERS

14




Lie on a bench with the feet flat on the ground and hold a dumbbell in the palms of both hands, with the thumbs surrounding the handle and arms extended:

- Inhale and lower the dumbbell behind the head, bending slightly at the elbows.
- Exhale and return to the initial position.


This exercise develops the bulk of the pectoralis major, long head of the triceps brachii, teres major, latissimus dorsi, serratus anterior, rhomboids, and pectoralis minor. The last three muscles stabilize the scapula so that the humerus can move from a stable base.

THE MOVEMENT


STABILIZERS OF THE SHOULDER BLADES



If you use this exercise to open the rib cage, you must work with light weights and avoid bending too much at the elbows. If possible, use a convex bench or place yourself across a horizontal bench and position the pelvis lower than the shoulder girdle. Take in a deep breath at the beginning of the movement and breathe out only at the end of the execution.




PERFORMING THE MOVEMENT AT A MACHINE





VARIATION POSITION ACROSS A BENCH

Placing yourself transversely across a bench opens the rib cage.


15 BARBELL PULLOVERS




THE MOVEMENT



INSERTIONS OF THE SERRATUS ANTERIOR



SERRATUS ANTERIOR MUSCLE



With arms extended, hold the barbell with an overhand grip and hands shoulder-width apart:


- Inhale and expand the chest as much as possible, lowering the barbell behind the head while bending slightly at the elbows.
- Exhale while returning to the initial position.

This exercise develops the pectoralis major, long head of the triceps brachii, teres major, latissimus dorsi, serratus anterior, rhomboids, and pectoralis minor.

This is an excellent movement for developing the flexibility and expansion of the rib cage. It should be performed with light weights using proper form and breathing.

4



BACK



| | | |
|----|--|-----|
| 1 | CHIN-UPS | 86 |
| 2 | REVERSE CHIN-UPS | 88 |
| 3 | STRETCHING THE LATISSIMUS DORSI AND TERES MAJOR | 89 |
| 4 | LAT PULL-DOWNS | 90 |
| 5 | BACK LAT PULL-DOWNS | 91 |
| 6 | TRICEPS BRACHII TEAR | 92 |
| 7 | CLOSE-GRIP LAT PULL-DOWNS | 93 |
| 8 | SEATED ROWS | 94 |
| 9 | WIDE GRIP SEATED ROWS | 95 |
| 10 | STRAIGHT-ARM LAT PULL-DOWNS | 96 |
| 11 | ONE-ARM DUMBBELL ROWS | 97 |
| 12 | DUMBBELL ROWS | 98 |
| 13 | BARBELL ROWS | 99 |
| 14 | T-BAR ROWS | 100 |
| 15 | T-BAR ROWS WITH ABDOMINAL SUPPORT | 101 |
| 16 | STIFF-LEGGED DEADLIFTS | 102 |
| 17 | SUMO DEADLIFTS | 103 |
| 18 | DEADLIFTS | 104 |
| 19 | BICEPS BRACHII TENDON TEAR | 106 |
| 20 | TRAP BAR DEADLIFTS | 108 |
| 21 | MUSCLES RECRUITED ACCORDING TO THE DIFFERENT TYPES OF DEADLIFT | 110 |
| 22 | LOW BACK PAIN SHOULD YOU ARCH YOUR BACK? | 111 |
| 23 | BACK EXTENSIONS | 112 |
| 24 | MACHINE BACK EXTENSIONS | 113 |
| 25 | STRETCHING THE BACK WITH A CHIN-UP BAR | 114 |
| 26 | UPRIGHT ROWS | 115 |
| 27 | BARBELL SHRUGS | 116 |
| 28 | DUMBBELL SHRUGS | 117 |
| 29 | TRAP BAR SHRUGS | 118 |
| 30 | MACHINE SHRUGS | 119 |
| 31 | STRETCHING THE DELTOIDS, TRAPEZIUS, AND NECK | 120 |
| 32 | STRETCHING THE TRAPEZIUS AND NECK | 121 |

1

CHIN-UPS



Hang from a fixed bar with a very wide overhand grip:

- Inhale and pull the chest up to the level of the bar.
- Exhale at the end of the movement.
- Return to the initial position with a controlled descent and begin again.

This exercise takes a certain amount of strength and is excellent for developing the latissimus dorsi and teres major and, when the shoulder blades come together at the top of the chin-up, the rhomboids and middle and lower portions of the trapezius. It also works the biceps brachii, brachialis, and brachioradialis.

Variations: If you stick out your chest, you can pull yourself up so the bar touches your chin. To increase the intensity, wear a weight belt. Keeping the elbows in next to the body during the movement contracts mainly the lateral fibers of the latissimus dorsi and develops the width of the back.


Bringing the elbows back and the chest out as you raise the chin to the bar mainly solicits the upper and central fibers of the latissimus dorsi and those of the teres major. This exercise develops the bulk of the back when the shoulder blades come together and the rhomboids and the upper and lower portions of the trapezius are used equally.

BRINGING THE BAR TO THE NECK WITH ARMS ALONGSIDE THE BODY



Bringing the elbows alongside the body mainly solicits the lateral fibers of the latissimus dorsi and develops the width of the back.

BRINGING THE BAR TO THE CHEST WITH ELBOWS BEHIND




Pulling the elbows back to raise the chin to the bar mainly solicits the upper and central fibers of the latissimus dorsi. This variation is excellent for developing the bulk of the back.

EVOLUTION

Originally, the teres major and latissimus dorsi muscles of our far-off ancestors played a role in their getting around on all fours by helping them to push off with the front paws.

As our ancestors became tree climbers, these muscles became powerful, specializing in vertical displacement. Returning to the ground, our more recent ancestors adopted bipedal displacement without losing the possibility of climbing. For this reason we possess powerful back muscles capable of pulling our bodies up, still allowing us to climb trees.


Comment: The main difference between our locomotor mechanism and that of our close simian relatives is the development of lower extremities that allow us to walk on two legs. Our chest and upper extremities have not developed differently and have the same structure and proportions. Contrary to popular opinion, apes do not have relatively big arms; we just have big legs!



In both monkeys and humans, the latissimus dorsi and teres major are particularly well developed.

2

REVERSE CHIN-UPS



Hang from a bar with an underhand grip, hands shoulder-width apart:

- Inhale and push out the chest as you raise the chin to the bar.
- Exhale at the end of the movement.


This movement develops the latissimus dorsi and teres major and is associated with the intense work of the biceps brachii and brachialis.

Therefore, it could be included in an arm workout program.

This exercise also contracts the middle and lower portions of the trapezius, the rhomboids, and the pectorals.

Performing this exercise takes a certain amount of strength; use a high pulley to make it easier.



MUSCLE INSERTIONS AT THE SHOULDER BLADE



POSTERIOR VIEW

ANTERIOR VIEW

STRETCHING THE LATISSIMUS DORSI AND TERES MAJOR





Stand with your legs slightly apart:

- Lean forward with the chest. With the arm extended, grasp a stable support such as a weightlifting machine or the frame of a squat cage.
- Place the palm of the other hand higher up on the support. With the arm extended, apply an increasingly powerful push against the machine while at the same time pulling with the other arm.

To accentuate the stretch of the latissimus dorsi and teres major, rotate the chest while trying to slowly raise the other shoulder.

When practiced regularly and incorporated into the first series of back-specific training, this stretch helps prevent tearing of the latissimus dorsi and teres major, which can occur during the execution of heavyweight pulls at the high pulley or tractions at the fixed bar with weight.

INSERTIONS OF LATISSIMUS DORSI AND TERES MAJOR ON THE HUMERUS (ANTERIOR VIEW)





ANTERIOR VIEW

The latissimus dorsi twists around the teres major to insert into the anterior surface of the humerus with a flat tendon.



3

LAT PULL-DOWNS




Sit facing the machine with the legs positioned under the pads, gripping the bar with a wide overhand grip: Inhale and pull the bar down to the sternal notch while puffing out the chest and pulling the elbows back. Exhale at the end of the movement.

This exercise develops the bulk of the back. It mainly works the upper and central fibers of the latissimus dorsi. The middle and lower portions of the trapezius, the rhomboids, the biceps brachii, the brachialis, and, to a lesser extent, the pectorals also contract.




BACK LAT PULL-DOWNS

4



ACTION OF TERES MAJOR AND LATISSIMUS DORSI



Sit facing the machine with the thighs positioned under the pads. Grasp the bar with a wide overhand grip:

- Inhale and pull the bar down to the back of the neck, bringing the elbows alongside the body.
- Exhale at the end of the movement.

This exercise develops the width of the back. It works the latissimus dorsi (mainly the lateral and lower fibers), the teres major, the forearm flexors (biceps brachii, brachialis, and brachioradialis), the rhomboids, and the lower portion of the trapezius. The latter two muscles come into play when the shoulder blades are pulled together. Back lat pull-downs help beginners develop enough strength to move on to chin-ups.




THE MOVEMENT



VARIATION AT A MACHINE WITH A FIXED AXIS



TRICEPS BRACHII TEAR



Although it is not the most-used muscle when working the back, the long head of the triceps brachii is the most frequently injured muscle during back lat pull-downs with heavy weights or during chin-ups with added weight.

The latissimus dorsi is a powerful, fan-shaped muscle that attaches the arm to the rib cage, and whose distal tendon is strongly attached to the humerus.

This is the main climbing muscle.

The long head of the triceps brachii, on the other hand, is a smaller muscle whose main function is to extend the forearm and secondarily to bring the arm toward the rib cage. In this way it complements the action of the latissimus dorsi.

Tearing of the long head of the triceps occurs when the muscle is fatigued, most frequently after an improper warm-up.

It only takes a sudden relaxation of the latissimus dorsi during chin-ups with added weight to immediately shift the tension to the long head of the triceps.


This tendon may partially tear, most often close to its insertion on the scapula. (Fortunately complete tears are infrequent.)

Unlike incapacitating shoulder injuries, which may completely halt upper-body training, a tear in the long head of the triceps is less devastating.

You can still perform back exercises such as seated rows or T-bar rows and movements for the triceps such as forearm extensions at a high pulley with the elbows next to the body despite the injury as long as you begin with lighter weights.

However, a brief rest period is recommended before beginning upper-body training.


Comment: Tearing the long head of the triceps may also occur during bench presses. To prevent this triceps tear, warm up with stretching exercises (see page 31).



TRICEPS STRETCH

CLOSE-GRIP LAT PULL-DOWNS

5



Sit and face the machine with knees positioned under the pads:

- Inhale and bring the handle to the sternum while expanding the chest and leaning slightly back with the torso.
- Exhale at the end of the movement.


This exercise develops the latissimus dorsi and teres major.

When the shoulder blades come together, the trapezius and the posterior deltoid contract.

As with every pulling exercise, the biceps brachii and brachialis contract, and when the palms face each other, the brachioradialis comes into play.

6

SEATED ROWS



Sit facing the machine, feet resting on the foot pad and the torso bent forward:


- Inhale and bring the handle to the base of the sternum by straightening the back and pulling the elbows back as far as possible.
- Exhale at the end of the movement and return smoothly to the initial position.

This exercise works the bulk of the back. It focuses the effort on the latissimus dorsi, teres major, posterior deltoid, biceps brachii, and brachioradialis, and, at the end of the movement when the shoulder blades come together, the trapezius and rhomboids.

While raising the chest, the spinal muscles (erector spinae) also contribute.


Allowing the weight to pull you on the return helps develop back flexibility.

Attention: To prevent back injury, never round the back when performing seated rows with heavy weights.



WIDE GRIP SEATED ROWS

7




Sit facing the machine with your feet on the foot pads and your chest flexed. Grasp the bar with an overhand grip (thumbs to the inside) wider than shoulder-width apart:

- Inhale and pull the bar to your chest, straightening the back and keeping your elbows raised.
- Exhale at the end of the movement and return smoothly to the initial position.

This is an excellent exercise for working the upper back behind the shoulders. The main muscles worked are the latissimus dorsi, teres major, posterior deltoid, infraspinatus, teres minor, arm flexors (biceps brachii, brachialis, brachioradialis), and, with the approximation of the shoulder blades, the rhomboids and the middle part of the trapezius. During the straightening of the chest, the erector spinae are also recruited.

Variation: Hold the bar in an underhand position (thumbs to the outside) to work the inferior part of the trapezius, rhomboids, and biceps brachii.





VARIATIONS WITH A WIDE GRIP

- 1 An underhand grip isolates the inferior portion of the trapezius.
- 2 An overhand grip isolates the posterior deltoid and the middle portion of the trapezius.


Attention: To prevent back injury, never round the back when performing seated rows with heavy weights.

8

STRAIGHT-ARM LAT PULL-DOWNS



MUSCLES USED




Stand and face the machine with feet slightly apart. Grip the bar with an overhand grip, arms extended and shoulder-width apart. Fix the back and contract the abdominal core:

- Inhale and bring the bar to the thighs, keeping the arms extended (elbows can be slightly bent).
- Exhale at the end of the movement.

This exercise, which works the latissimus dorsi, strengthens the teres major and the long head of the triceps, which stabilizes the arm-trunk hinge.


Comment: Many swim coaches use this exercise to develop a powerful crawl stroke.



THE MOVEMENT

ONE-ARM DUMBBELL ROWS

9




Grasp a barbell with the palm facing in; use the opposite hand and knee on the bench to support the back:

- Inhale and lift the upper arm and elbow as high as possible next to the body with the elbow bent.
- Exhale at the end of the movement.


To maximize the contraction, rotate the torso slightly toward the working side at the end of the row.

This exercise mainly works the latissimus dorsi, teres major, and posterior deltoid, and, at the end of the contraction, the trapezius and rhomboids. The forearm flexors (biceps brachii, brachialis, and brachioradialis) are also used.



10

DUMBBELL ROWS



THE MOVEMENT

Stand with the legs slightly bent and the chest leaning forward at approximately a 45-degree angle. Keep your back very flat and hang your arms by your sides; hold a dumbbell in each hand with an overhand grip, palms facing inward:

- Inhale and isometrically contract the abdominal core. Pull the dumbbells as high as possible, keeping the elbows close to the body. Squeeze the shoulder blades together at the end of the movement.
- Return to the initial position and exhale.

This exercise recruits the latissimus dorsi, teres major, posterior deltoid, forearm flexors (biceps brachii, brachialis, brachioradialis), and, with approximation of the shoulder blades, the rhomboids and trapezius.

The tilted position of the chest works the spinal muscles isometrically.

By varying the angle of the chest, it is possible to focus the work onto specific parts of the back:



1. Keeping the chest up mainly works the superior trapezius.
2. Keeping the chest closer to horizontal works the dorsals, teres major, rhomboids, and middle and inferior trapezius.




Attention: To avoid injury, never round the back during execution of the movement.

BARBELL ROWS


11



THE MOVEMENT



1 Hands in pronation (overhand grip) emphasizes the rhomboids and lower portion of the trapezius.



2 Hands in supination (underhand grip) emphasizes the upper portion of the trapezius and the biceps brachii.

Stand with legs slightly bent and grasp the bar with an overhand grip with the hands wider than shoulder-width apart. With the back straight, lean forward at the waist 45 degrees, so that the bar is at knee level:

- Inhale and hold the breath during an isometric contraction of the core and pull the barbell up to the chest.
- Return to the initial position and exhale.

This exercise contracts the latissimus dorsi, teres major, posterior deltoid, and the forearm flexors (biceps brachii, brachialis, brachioradialis). When the shoulder blades come together, the rhomboids and trapezius contract.

The forward lean of the torso uses the spinal muscles in an isometric contraction.


Varying the hand position in width and grip (overhand or underhand) along with changing the angle of the torso allows you to work the back from a variety of angles.

Attention: To prevent injury, never round the back during this exercise.



12

T-BAR ROWS



Straddle the bar with the legs slightly bent and lean forward at the waist about 45 degrees with a flat back. Grasp the bar with an overhand grip:

- Inhale and raise the bar to the chest.
- Exhale at the end of the movement.


This exercise is similar to bent rows and allows you to concentrate on working your back because you do not have to focus too much effort on positioning.

This exercise uses mainly the latissimus dorsi, teres major, infrapinatus, rhomboids, trapezius (mainly the middle portion), and the flexors of the forearm.

The forward lean isolates the abdominal and spinal muscles in isometric contraction.

Using a supinated (underhand) grip transfers some of the effort to the biceps brachii and the upper portion of the trapezius at the end of the pull.




Some machines are equipped with parallel handles that allow a grip between pronated and supinated, which contracts the brachioradialis more intensely.



T-bar rows also work the forearm flexors.

T-BAR ROWS WITH ABDOMINAL SUPPORT

13



VARIATION AT A MACHINE
REPRODUCING THE MOVEMENT
AS PERFORMED AT A T-BAR

Rest against the incline bench:

- Inhale and bring the bar to the chest with an overhand grip.
- Exhale at the end of the movement.


This exercise is similar to bent rows and allows you to concentrate on working your back because you do not have to focus too much effort on positioning.

It mainly uses the latissimus dorsi, teres major, posterior deltoid, arm flexors, trapezius, and rhomboids.

Some machines are equipped with an abdominal support, which eliminates the work of the abdominal and spinal muscles. However, when using heavy weights, the rib cage is compressed against the abdominal-support pad, which interferes with breathing and makes the exercise painful to perform.

Comment: A pronated (overhand) hold shifts some of the effort to the biceps brachii and the upper portion of the trapezius at the end of the pull.

14 STIFF-LEGGED DEADLIFTS



Stand with the feet slightly apart, facing the bar as it rests on the ground:

- Inhale and bend forward at the waist with the chest forward, back arched, and legs as straight as possible.
- Grasp the bar with an overhand grip. Keeping the arms relaxed, stand up straight by rotating the hips. Keep the abdominal muscles tight and a slight arch in the back for support.
- Exhale at the end of the movement.
- Bend forward and return to the initial position, but without returning the bar to the floor.


To avoid injury, keep the back straight.

This exercise contracts the deep spinal muscles on either side of the spinal column that straighten the spine. Straightening the torso by tilting the pelvis from front to back contracts the gluteus maximus and hamstrings (except the short head of the biceps femoris).

Deadlifting from the ground with extended knees stretches the back of the thighs.

To increase the intensity, stand on a box so that the feet are higher than the bar on the ground.




Comment: To stretch the hamstrings, perform the stiff-legged deadlift with very light weights. The greater the weight, the more the gluteal muscles take over from the hamstrings to straighten the pelvis to vertical.



SUMO DEADLIFTS

15

THE MOVEMENT

DEEP MUSCLES OF THE BACK
USED DURING THE SUMO DEADLIFT

Stand facing the bar, with legs wider than shoulder-width apart and toes pointing out in line with the knees:

- Inhale and bend the legs until the thighs are horizontal to the ground; grasp the bar with an overhand grip about shoulder-width apart. If you are lifting very heavy weights, use a reverse grip (grasp the bar with one overhand and one underhand grip) to keep the bar from rolling.
- Hold your breath and contract the core, slightly round the back, and extend the legs, bringing the torso vertical and pulling the shoulders back.
- Exhale at the end of the movement.
- Return the bar to the ground while holding your breath. Never round your back.



The difference between this and the classic deadlift is that this exercise works the quadriceps and adductor muscles intensely. Because the pelvis is not as tilted, it works the back less.

Comments: When beginning the movement, slide the bar along the shins. High reps (10 maximum) with light weights strengthen the lumbar region and work the thighs and the gluteal muscles.


When using heavy weights, perform this exercise with great caution to prevent injuries to the hip joints, adductor group of the thighs, and the lumbosacral junction. The sumo deadlift is one of the three powerlifting movements.

16

DEADLIFTS



With vertebral flexion, the disc is compressed in front and expanded in back. The fluid of the nucleus pulposus migrates to the back and may compress on the neural elements (which causes sciatic pain).




Vertebral foramen (hole through which a spinal cord nerve runs)

Comment: By limiting the rolling of the bar, the reverse grip allows you to lift much heavier weights.

Reverse power grip

Classic grip







Stand facing the barbell, legs slightly apart with the abdominal muscles contracted and the back slightly arched. Bend the knees until the thighs are almost parallel to the floor. This position will vary depending on your physical structure and the flexibility of your ankles. (The thighs will be horizontal for someone with short thigh bones and arms. The thighs will be a little above the knees for someone with long thigh bones and arms.) Take an overhand grip on the bar with your hands slightly more than shoulder-width apart. You can also use an over-under grip (one palm faces forward and the other faces back) to prevent the bar from rolling and to work with much heavier weight.

- Inhale, hold the breath, contract the abdominal and low back muscles, and lift the bar by straightening your legs and allowing the bar to slide up the shins.
- When the bar reaches the knees, extend your torso while straightening your legs so you are standing erect with your arms straight down at your sides, exhaling as you complete the movement.
- Hold this straightened position for 2 seconds, then return the weight to the floor, making sure you do not hyperextend or arch your back.

Throughout the exercise, keep your back straight.

This exercise works nearly every muscle in the body and is effective for developing the lumbosacral and trapezius muscles. It also works the gluteal muscles and quadriceps intensely.

The deadlift, along with the bench press and the squat, make up the exercises in powerlifting competitions.



MUSCLES USED DURING THE DEADLIFT (HIGHLIGHTED)


Attention: No matter what the exercise, as soon as heavy weights are involved, it is essential to create a *block*:

1. Expanding the chest and holding a deep breath fills the lungs, which supports the rib cage and prevents the chest from collapsing forward.
2. Contracting the abdominal muscle group supports the core and increases the intra-abdominal pressure, which prevents the torso from collapsing forward.
3. Finally, arching the low back by contracting the lumbar muscles positions the spinal column in extension.

These three actions together are referred to as *blocking*, which keeps you from rounding the back (vertebral flexion). A rounded back when lifting heavy weights can cause a herniated disc.



BICEPS BRACHII TENDON TEAR



Tearing the long head of the biceps brachii is by far the most common serious sport-related biceps injury.

Generally, it occurs in a muscle, already weakened by tendinitis, after a sudden backward movement of the arm (e.g., during a throw). This movement is relatively common in baseball, tennis, and any sport involving a throwing action, but it also occurs in the snatch in weightlifting. During this motion, tension is suddenly placed on the long head of the biceps brachii, most often where its tendon passes through the bicipital groove of the humerus.

Weightlifting, specifically the deadlift, with heavy weights can cause another characteristic biceps brachii injury.

A common practice when using heavy weights in the deadlift is to use a reverse grip (one overhand grip and one underhand grip) to prevent the bar from rolling in the hands.

This technique, although usually safe, can in rare instances cause the tearing or the pulling away of the inferior tendon of the biceps brachii where the muscle inserts onto the humerus.

During the positive phase of the deadlift, the effort is mainly exerted by the leg, gluteal, back, and abdominal muscles. The arms hang down, completely extended and relaxed. Unfortunately, the slight shortening caused by contracting either head of the biceps brings the hand into supination (the biceps being the strongest supinator), which with extra heavy weights may cause complete rupture of the tendon at the radius. This injury occurs at the distal attachment because as the arms hang next to the body, the proximal tension is divided between the short and long heads of the biceps brachii, whereas distally, only one tendinous insertion supports the tension.

Compared to other tendon tears such as the pectoralis major or the adductors of the thigh, in which the pain is unbearable and stops the athlete from continuing, the pain of a biceps tendon tear is relatively mild despite the seriousness of the actual injury. In competitive powerlifting, athletes have continued their lift despite the biceps tendon tear incurred during that lift.


After the accident the diagnosis is obvious: Swelling caused by hemorrhaging appears in the forearm. But what is most striking is the appearance of the biceps

brachii, which becomes ball shaped at the upper arm close to the pectoralis major and the deltoid, revealing the brachialis muscle lower down.

Despite the tear, the brachialis, brachioradialis, extensor carpi radialis longus and brevis, and pronator teres muscles can still flex the arm, just not as strongly. Supination of the forearm becomes much more of a problem because the end range of this movement relies only on the supinator muscle.


If this injury is not immediately treated with surgery to reattach the biceps tendon onto the radius, irreversible retraction of the muscle will occur with fibrous change. And although moving the arm will still be possible, there will be permanent loss of strength in flexion and supination. It is possible to prevent this injury by regularly working the biceps, not to develop the muscle, but to strengthen its tendon. For this reason add forearm flexion isolations using a bar in a series of "cheats" by leaning the chest back to give the bar a boost. If practiced regularly, this technique reinforces the distal tendon of the biceps by the tension it places on it. Nevertheless it must be performed carefully without rounding the back to avoid injury.

Pectoralis major
Deltoid
**Biceps brachii
(sectioned and retracted)**
Triceps brachii
Brachialis



**BICEPS BRACHII MUSCLE
RETRACTED WITH TEARING
OF ITS DISTAL TENDON**

NORMAL BICEPS BRACHII MUSCLE



**TYPICAL APPEARANCE OF AN UNTREATED
DISTAL BICEPS TENDON TEAR**


If, after tearing the distal tendon of the biceps brachii, surgery to reattach it to the radius is not performed quickly, permanent retraction and atrophy of the muscle will occur.



Biceps tendon on the arm of the supinated hand can tear during a heavy deadlift.


17

TRAP BAR DEADLIFTS



Stand well centered in the bar (be careful because poor centering will lead to lateral instability). Legs are slightly apart and back is well fixed and slightly bent:

- Flex the legs in order to bring the thighs close to horizontal; this position can vary depending on the flexibility of the ankles and the individual morphology (for example, for people with short femurs and arms, the thighs will be horizontal; for people with long femurs and arms, the thighs will be more or less higher than horizontal).
- With arms extended, grasp the handles of the bar, carefully centering the grip. (Attention: With heavy weights on a trap bar, a badly adjusted grip will push the bar forward or back.)
- Inhale, contract the abdominal girdle and the lumbar region, and raise the bar by straightening the legs without ever rounding the low back. Exhale at the end of the effort.
- Maintain the extension of the body for 2 seconds, then replace the bar while keeping the abdominal girdle and the lumbar region contracted.




As in the classic deadlift (page 104), the deadlift with a trap bar works all the muscles of the body, but the centered position of the bar allows you to decrease the lean of the torso, which will limit the intensity of the work on the lumbar region of the gluteals by throwing some of the effort onto the quadriceps. Thus this movement may be included in a specific program of thigh work and in some cases can replace the squat.

With heavy weights, the superior part of the trapezius is strongly recruited.


Comment: If you have low back pain, this exercise is safer than the classic deadlift.



MUSCLES RECRUITED ACCORDING TO THE DIFFERENT TYPES OF DEADLIFT



ANGLE OF THE CHEST ACCORDING TO THE DIFFERENT TYPES OF DEADLIFT






LOW BACK PAIN

Back pain is the most common problem of the lumbar spine region.

Generally, it is not serious and is most often caused by the shortening of the small, deep vertebral muscles that attach to the transverse processes.

If, during a poorly executed rotation or extension of the spine, one of these muscles is overstretched or is torn, it will automatically shorten along with its neighboring muscles and the superficial erector spinae. The back muscles cramp in pain; however, this cramping limits movement that otherwise might tear or increase the tearing of the small deep muscle.

This general shortening of a portion of the back muscles often disappears when the small deep muscle heals. But sometimes the back pain becomes entrenched, and even after the muscles heal, the local shortening can last several weeks and in some people for years.



SMALL DEEP MUSCLES OF THE BACK


Comment: Although not serious in and of itself, lumbago, which is a painful contracture of the back muscles, can be part of more serious vertebral injuries such as herniated discs, tears in the paravertebral muscles and ligaments, and fractures.




SHOULD YOU ARCH YOUR BACK?

For people without vertebral problems, arching the back during an exercise is not risky. In fact, with movements such as the squat (page 126) or the deadlift (page 104), where the back tends to round, arching the back can prevent injury. However, for some people arching the back during an exercise can be very dangerous.

- For people suffering from congenital spondylolysis (incomplete fusing of the vertebral arch), putting the lumbar spine in extension can cause the vertebra to slide (spondylolisthesis), which may cause serious nerve compression and lead to sciatica.
- For people who are not fully grown or people experiencing osteoporosis, extending the lumbar spine may lead to spondylolysis because of fractures in the vertebral arch. This fracture in the posterior anchoring system of the vertebra may allow the vertebra to slide forward and seriously compress the neural elements (which leads to sciatica).







LUMBAR VERTEBRA FROM ABOVE



With spondylolysis (fracture of the vertebral arch) the vertebra may slide forward (spondylolisthesis) and compress the neural elements, causing sciatica.

18 BACK EXTENSIONS



Lie facedown on a Roman chair and place the ankles under the roller pads. Because the axis of flexion passes through the coxo-femoral joints, the pubic bone should not rest on the support pad:

- With the torso bent forward, extend the back to horizontal.
- Raise the head and continue into hyperextension by arching the lumbar spine. This must be performed carefully to protect your low back.

This exercise mainly develops the group of paraspinal erectors of the spine (iliocostales, longissimus thoracis, spinalis thoracis, splenius, and semispinalis capitis) and quadratus lumborum and, to a lesser degree, the gluteus maximus and the hamstrings, except for the short head of the biceps femoris. Complete flexion of the torso develops the flexibility of the lumbosacral mass. Supporting the pelvis on the bench, so that the axis is displaced to the back of the body, focuses the movement completely at the lumbosacral level but less intensely, given the range of motion and the greater power of the lever arm.


To increase the intensity, sustain the horizontal position of the torso at the end of the extension for a few seconds. Using an incline bench makes this exercise easier for beginners to execute.

Variations

- Performing the torso extension with a bar on the shoulders stabilizes the upper back, which focuses the effort on the lower part of the erector spinae muscles.
- The back extension machine allows you to focus on the lumbosacral mass of the spinal muscles (see page 113, machine back extensions).
- To increase the intensity, perform the exercise while holding a weight to the chest or behind the neck.

MACHINE BACK EXTENSIONS

19



Sit at the machine, with the torso leaning forward and the pad of the machine at shoulder-blade (scapula) level:

- Inhale and press back, straightening the torso as much as possible.
- Return slowly to the initial position while exhaling and begin again.

This exercise works the erector muscles of the spine, focusing the effort on the low back, specifically the lumbosacral mass of the spinal muscles.

This exercise is excellent for beginners. Done in sets of 10 to 12 repetitions, it develops the strength to progress to more technically demanding exercises for the back.


To perform this exercise with heavier weights, reduce the number of repetitions in the set.

Because the machine regulates the range of motion and the weight, the number of repetitions may vary during the same session.

Example: Two series of 15 repetitions with moderate weights and complete range of performance followed by two series of 7 repetitions with more weights and reduced range.




STRETCHING THE BACK WITH A CHIN-UP BAR



Bend head forward and try to touch chin to chest.

Slowly release the back to feel the stretch on the small intervertebral muscles.

DIAGRAM OF VERTEBRAE



1 When you lift heavy weights, such as during squats or in deadlift exercises, the intervertebral discs can become pinched and the nucleus can migrate to the periphery.

2 When you hang from a chin-up bar, the small intervertebral ligaments and muscles stretch, the intervertebral disc compression decreases, and the nucleus pulposus can return to its position at the center of the disc.

Hang from a chin-up bar with a wide overhand grip (thumbs facing each other):

- Inhale and exhale slowly, concentrating on the relaxation of your body, which allows the back muscles to relax and the pressure inside the discs to equilibrate as well as relax the small paravertebral muscles (interconnecting the vertebrae), which are often painfully contracted.
- When you are relaxed, lean your head forward, trying to touch your sternum with your chin. This will stretch the upper and middle back.

To accentuate the stretch, swing gently or ask a partner to grasp your hips on each side and slowly pull down.

This stretch is fundamental. When practiced regularly at the end of squat and deadlift sessions (or any other heavyweight exercises that would have compressed the spine), it helps over time to limit the deterioration of the intervertebral discs and reduces the risk of disc herniation (see page 134).


Variation: If you grip strongly with the hands, the latissimus dorsi and teres major muscles are stretched more intensely.

Comment: During this exercise, you often hear cracking of the spine followed by a pleasant feeling of freedom and relaxation of the vertebral column.

These cracking sounds are caused by the release of the paravertebral muscles acting like a bellows that, when opened, decompresses the small intervertebral and costovertebral articulations.

UPRIGHT ROWS

20




Stand with the legs slightly apart, keeping the back straight and grasping the barbell with an overhand grip. The grip should be hand width or slightly wider:

- Inhale and pull the barbell up along the front of the body to the chin, raising the elbows as high as possible.
- Exhale and lower the barbell with a controlled movement.



This exercise mainly uses the superior portion of the trapezius as well as the deltoids, levator scapulae, biceps brachii, brachialis, forearm muscles, abdominal muscles, gluteal muscles, and lumbosacral group.

A wider grip uses the deltoid more than the trapezius.



21

BARBELL SHRUGS



TRAPEZIUS

Stand with the legs slightly apart and face the bar that is either on the ground or on a stand:




- Grasp the bar with an overhand grip with the hands slightly wider than shoulder-width apart, or use a reverse grip if the weight is heavy.
- Shrug the shoulders while keeping the arms relaxed, back straight, and abdominal muscles contracted.

This exercise develops the upper portion of the trapezius, mainly its occipital-clavicular fibers, and the levator scapulae.

Comment: Use a reverse grip when working with heavy weights. To balance the work on the trapezius, change the hands with each set. For example, perform one set with the right hand overhand and the left hand underhand, then switch on the following set.

DUMBBELL SHRUGS

22





INITIAL POSITION

Stand with the legs slightly apart, head upright or slightly flexed forward, and arms relaxed at the sides. Grip a dumbbell in each hand:

- Shrug the shoulders with an anterior to posterior rotation.
- Return to the initial position.

This exercise contracts the upper, or clavicular, and middle portions of the trapezius, the levator scapulae, and the rhomboids when squeezing the shoulder blades together and rotating the shoulders to the back.

TRAPEZIUS ACTION




EXECUTING THE ROTATION AT THE END OF THE MOVEMENT

Comment: It is impossible to rotate the shoulders when using heavy weights.

23

TRAP BAR SHRUGS



INITIAL POSITION

Stand with your legs slightly apart. Face the bar on the floor or on a support:

- Grasp the bar, making sure that the hands are centered. (**Attention:** When you have heavy weights on a trap bar, gripping the bar without adjusting the hands properly will cause the bar to swing forward or backward.)
- With the head straight or slightly forward, arms relaxed, back very straight, and abdominals contracted, perform shoulder shrugs.

This exercise mainly works the superior portion of the trapezius, which inserts onto the clavicle, acromion, and spine of the scapula and ascends into the superior nuchal line of the cranium.



On a deeper level, the minor and major rhomboids and levator scapulae are also solicited.

As the name indicates, the trap bar was initially created to work the trapezius muscles; it allows for lifting heavier weights without scraping the thighs as with straight bars or dumbbells.

Comment: People with long clavicles will find it more difficult than people with shorter clavicles to perform shoulder shrugs with heavy weights.

MACHINE SHRUGS

24






Comment: Just as the road bed of a bridge is suspended by metal cables, the clavicles and shoulder blades are suspended by the trapezius.

Splenius
Sternocleidomastoid
Levator scapulae
Scalenes

Spine of scapula
Infraspinatus
Trapezius, lower portion
Teres major
Latissimus dorsi
Anconeus
Extensor digitorum

Trapezius, upper portion

Deltoid
Teres minor
Triceps brachii
Brachioradialis
Extensor carpi radialis longus
Extensor carpi radialis brevis



**FINAL POSITION
TRAPEZIUS CONTRACTED**



Stand facing the machine and hold the handles with an overhand grip slightly wider than shoulder width, or if the machine allows, with the palms facing each other:

- Shrug the shoulders, keeping the head and back straight.

This exercise allows you to perform long sets. It is excellent for developing the upper portion of the trapezius and the levator scapulae.



STRETCHING THE DELTOIDS, TRAPEZIUS, AND NECK




MUSCLES STRETCHED

Stand with your legs slightly apart and back very flat. Place one arm behind your back and clasp that wrist with your other hand. Pull the arm slowly down and toward the outside of your body to feel the stretch of the deltoid (mainly the posterior and middle portions) as well as the trapezius.

Variation: To better feel the stretch at the neck, perform this exercise while slowly bending the head to the side opposite of the shoulder being stretched.

This variation allows you to stretch the deep complex muscles at the edge of the cervical spine as well as the scalene and sternocleidomastoid.

STRETCHING THE TRAPEZIUS AND NECK



Place one hand over your head and pull gently, bending the head to the side. This exercise stretches the sternocleidomastoid, scalene group, superior portion of the trapezius, splenius capitis, splenius cervicis, and deeper in the semispinalis capitis as well as the small muscles of the spine such as the longus colli, rectus capitis anterior, rectus capitis lateralis, and longus capitis.



Attention: Always perform this exercise progressively, pulling carefully on the head.



Comment: To better feel the stretch on the superior portion of the trapezius, lower the shoulder at the same time.



ABOUT THE NECK

In quadrupeds as well as the anthropomorphic ape, such as the gorilla, the muscles of the neck are particularly powerful and developed to maintain the head and prevent it from dropping forward.



But in humans, who have moved to complete bipedal uprightness with the head perched at the top of the spine, the muscles of the neck no longer serve to powerfully hold the head up but more to subtly balance the head on the spine.



1 In human evolution, the smaller face and the development of the bipedal stance and encephalon has caused the occipital foramen to move toward the center of the cranial base. The head positioned at the top of the spine requires the muscles of the neck to play only a subtle stabilizing role.

2 In the gorilla, a partial quadruped with a bigger face along with the posterior position of the occipital foramen, the muscles of the neck are particularly well developed and powerful to prevent the head from falling forward.



5 LEGS




| | |
|---|------------|
| 1 DUMBBELL SQUATS | 123 |
| 2 SQUATS WITH A DUMBBELL HELD BETWEEN THE LEGS | 124 |
| 3 FRONT SQUATS | 125 |
| 4 SQUATS | 126 |
| SQUAT-SPECIFIC STRETCH | 128 |
| 5 POWER SQUATS | 129 |
| ADAPTING TRAINING TO YOUR MORPHOLOGY | 130 |
| DISC HERNIATION | 134 |
| 6 INCLINE LEG PRESSES | 135 |
| 7 HACK SQUATS | 136 |
| 8 BOX SQUATS | 137 |
| 9 LEG EXTENSIONS | 138 |
| STRETCHING THE QUADRICEPS | 139 |
| 10 LYING LEG CURLS | 140 |
| 11 STANDING LEG CURLS | 141 |
| 12 SEATED LEG CURLS | 142 |
| HAMSTRING MUSCLE TEARS | 143 |
| 13 GOOD MORNINGS | 144 |
| STRETCHING THE HAMSTRINGS | 145 |
| 14 CABLE ADDUCTIONS | 146 |
| 15 MACHINE ADDUCTIONS | 147 |
| 16 FLEXION AND EXTENSION OF THE FEET | 148 |
| 17 STANDING CALF RAISES | 149 |
| 18 ONE-LEG TOE RAISES | 150 |
| 19 DONKEY CALF RAISES | 151 |
| 20 SEATED MACHINE CALF RAISES | 152 |
| 21 SEATED BARBELL CALF RAISES | 153 |
| STRETCHING THE CALF | 154 |
| SHORT CALF, LONG CALF | 154 |

DUMBBELL SQUATS

1



QUADRICEPS MUSCLE



Stand with the feet slightly apart, a dumbbell in each hand and the arms relaxed, looking straight ahead:

- Inhale, slightly arch the back, and bend the knees.
- When the thighs reach horizontal, extend the legs to return to the initial position.
- Exhale at the end of the effort.

This exercise mainly works the quadriceps and the gluteal muscles.



INITIAL POSITION


Comment: There is no point in working with heavy weights. Working with moderate weights in sets of 10 to 15 repetitions provides the best results.

2

SQUATS WITH A DUMBBELL HELD BETWEEN THE LEGS



INITIAL POSITION



Stand with your legs apart and toes pointing out. Hold a dumbbell in your hands:

- Look straight ahead, arch your back slightly, inhale, and bend your knees.
- When the thighs reach horizontal, extend your legs to return to the initial position.
- Exhale at the end of the movement.

This exercise works the quadriceps as well as the gluteal muscles.

Comment: The wide-leg position works the adductors.

FRONT SQUATS


3



THE MOVEMENT



Lowering the thighs by bending the knees while the legs are held by equipment focuses a major part of the work on the quadriceps muscles.



Stand with the legs about shoulder-width apart and hold the bar with an overhand grip as it rests on the upper pectoral muscles and the anterior deltoid:

- Inhale deeply to maintain intrathoracic pressure, which prevents the torso from collapsing forward, slightly arch the low back, contract the abdominal core, and bend the knees to lower the thighs horizontal to the floor.
- Return to the initial position and exhale at the end of the movement.

Stick out the chest and raise the elbows as high as possible to prevent the barbell from sliding forward.

Even though the barbell is in front, keep the back upright and don't lean the torso forward. To make the exercise easier, place blocks under the heels.

This type of squat focuses a greater part of the effort onto the quadriceps and is performed with lighter weights than the classic squat. This exercise also contracts the gluteal muscles, hamstrings, abdominal core, and the erector spinae. This is a movement frequently used in weight training because it corresponds perfectly with the work the thighs do at the end of a snatch.



CORRECT POSITION




INCORRECT POSITION



4

SQUATS



The squat is the number one bodybuilding movement: It uses nearly the entire muscular system, and it also works the cardiovascular system. It helps develop thoracic expansion, and therefore, respiratory capacity:


- With the barbell resting on a stand, slide under the bar and place it on the trapezius a bit higher than the posterior deltoid. Grasp the bar firmly with the hands at a comfortable width and the elbows back.
- Inhale deeply (to maintain the intrathoracic pressure, which will prevent the torso from collapsing forward), slightly arch the back by rotating the pelvis forward, contract the abdominal core, look straight ahead, and remove the barbell from the stand.
- Step back one or two steps and stop with both feet parallel to each other (or toes pointing slightly outward) and about shoulder-width apart. Bend forward from the hips (the axis of flexion should pass through the coxofemoral joints) and avoid rounding the back in order to prevent injury.

- When the thighs are horizontal to the floor, straighten the legs and lift the torso to return to the initial position.
- Exhale at the end of the movement.

The squat mainly works the quadriceps, gluteal muscles, adductor group, erector spinae, abdominal muscles, and the hamstrings.

Comment: The squat is one of the best exercises for developing the shape of the buttocks.

TORSO ANGLE BASED ON PHYSICAL DIFFERENCES





Variations

1. People with rigid ankles or long femurs can place a block under their heels to keep from tilting the torso too much. This variation isolates the quadriceps.
2. Lowering the bar onto the posterior deltoid increases the leverage of the back, which helps you lift heavier weights. This technique is essential for powerlifters.
3. The squat can be performed at a frame, which keeps you from tilting the torso and lets you focus on the quadriceps.

FOOT PLACEMENT IN THE SQUAT

When executing the classic squat, that is, with the feet approximately shoulder-width apart, you must place the feet properly. They should be parallel or slightly pointed to the outside. However, you must take into consideration your unique physical structure and make adjustments as necessary to ensure that the feet are in line with the knees.

Example: If you naturally walk with the feet pointed out, perform the squat with the feet pointed out.



1 CORRECT POSITIONS


When executing squats, hold the back straight.

Given the variations in each person's physical structure (different leg lengths and ankle flexibility) and the variations in technique (width of stance, use of heel blocks, barbell higher or lower), the tilt of the torso will vary; however, the lean should start at the hips.

2 INCORRECT POSITION

Never round the back when executing squats.

This mistake is responsible for most lumbar spine injuries, especially herniated discs.



TRADITIONAL HORIZONTAL SQUAT



COMPLETE SQUAT

Attention: To feel the working of the gluteal muscles, lower the thighs to horizontal.

1-2-3: NEGATIVE PHASE

4: COMPLETE SQUAT


To feel the working of the gluteal muscles even more, lower the thighs past horizontal. However, this technique can only be performed by people with flexible ankles or short femurs. Furthermore, you must perform the complete squat carefully and avoid the tendency to round the low back, which can lead to serious injury.

Attention: No matter what the exercise, as soon as heavy weights are involved, it is essential to create a *block*.

1. Expanding the chest and holding a deep breath fills the lungs, which supports the rib cage and prevents the chest from collapsing forward.
 2. Contracting the abdominal muscle group supports the core and increases the intra-abdominal pressure, which prevents the torso from collapsing forward.
 3. Finally, arching the low back by contracting the lumbar muscles positions the spinal column in extension.
- These three actions together are referred to as *blocking*, which keeps you from rounding the back (vertebral flexion). A rounded back when lifting heavy weights can cause a herniated disc.



SQUAT-SPECIFIC STRETCH




To avoid tearing muscles when practicing the squat, do some stretching exercises at the beginning of the workout while warming up and between the first series. One stretching exercise most often performed by powerlifters involves crouching deeply while holding onto a stable support such as the post of a weight training machine. This move, corresponding perfectly to the flexion of a squat, allows the adductors to be stretched favorably, especially the adductor magnus; this muscle is frequently injured during excessive tilting of the torso with heavy weights. The quadriceps (except for the rectus femoris), gluteus maximus, and the deep external rotator muscles of the hip stabilize and also slow down the anterior tilting in the crouch position.

Comment: To really feel the stretch on the inside of the leg, you can shift your body weight from one leg to the other.

POWER SQUATS

5




This movement is performed the same as the classic squat, except that the legs are farther apart and the toes point out, which works the inner thigh intensely.

The working muscles are as follows:

- Quadriceps
- Adductor muscle group (adductor magnus, adductor longus, adductor brevis, adductor pectineus, and gracilis)
- Gluteal muscles
- Hamstrings
- Abdominal muscles
- Lumbosacral muscle group

Comment: In the squat with legs farther apart, the torso is more upright at the bottom of the squat than in the classic squat. Some powerlifters choose the wide-leg technique in order to limit the stress on the back. However, certain powerlifters using heavy weights prefer the classic squat for relieving pressure on the low back because the torso can rest on the thighs at the bottom of the movement.





ADAPTING TRAINING TO YOUR MORPHOLOGY

Short-Limbed and Long-Limbed People

In weightlifting, it is important to consider your individual morphological type in your choice of form for exercises, particularly the squat and deadlift. Both these exercises work the muscles quite differently in short-limbed and long-limbed people.

A short-limbed person is someone with a proportionately longer torso and shorter limbs, whereas a long-limbed person has a relatively shorter torso and longer extremities. This has nothing to do with waistline, muscular development, or fat (a person can be short and long limbed or tall, thin, and short limbed).

Short-limbed people will be more adept at performing a squat. In fact, because of the shortness of the femur, the torso will tilt only a little, limiting the stress on the low back and the hamstring muscles, allowing the movement to be performed in relative security, and focusing almost completely on the quadriceps. It is therefore not surprising that almost all squat champions are in this morphological category. Extreme examples are the dwarves monopolizing the podiums in the small categories in powerlifting.


Long-limbed people will have more difficulty performing the squat. In fact, because of the length of the femurs, the torso will lean forward dramatically. This will place a dangerous amount of tension on the hamstring muscles as well as the adductor magnus and the gracilis. Long-limbed people constantly need to struggle to avoid training in an anterior imbalance.

People with long limbs will also have to concentrate on the position of the back to avoid rounding it, a fault that can lead to severe vertebral injuries—that is, disc herniation.


This type of long-limbed squat intensely works the gluteus maximus muscles (which come into play when straightening the pelvis and thus the chest) as well as the erector spinae muscles, which struggle against the rounding of the back.

The squat is excellent for developing powerful gluteals and good lumbar muscles in long-limbed people, but they will need to concentrate a great deal on their positioning.

1 SHORT-LIMBED TYPE




2 LONG-LIMBED TYPE




TILT OF THE TORSO IN THE SQUAT ACCORDING TO THE DIFFERENT MORPHOLOGIES

1 Long-limbed type:
Long legs
Short torso
Extremely tilted torso
Significant cantilever



2 Extremely short-limbed type
(dwarfism, achondroplasia):
Short legs
Long torso
Very little tilt of the torso
Less cantilever



Furthermore, the squat exercise becomes dangerous as soon as heavier weights are added. Long-limbed people who wish to target the work onto the quadriceps should work with machines, especially the hack squat (page 136).

Short-limbed people dominate squat competitions, but they don't do as well in deadlift competitions. Because of their shorter extremities, they are forced to flex the legs while grasping the bar on the floor. They must bring the femurs practically horizontal. Initiating the movement in this position requires an enormous amount of energy.

Long-limbed people initiate the deadlift with legs partially bent, a position in which the quadriceps can push the most. Despite a greater forward tilt of the back and more intense work on the gluteals and the erector spinae muscles, their morphology allows them to lift significantly heavier weights than short-limbed people can. This is why long-limbed people are essentially the champions of the deadlift.

Flexibility of the Ankles

Flexibility of the ankles has a big influence on the execution of the squat.

Whether it is due to a musculotendinous restriction (such as in retraction of the calves) or bone restriction, if dorsiflexion (raising the front part of the foot) is limited, the squat technique will be profoundly altered.


In fact, the lack of flexibility at the ankles will limit the forward tilt of the tibias and therefore the anterior movement of the knees. This forces the squat to be performed with the gluteals very far back and the back very tilted forward, which works the gluteals and erector spinae intensely.

An excessive tilt of the torso caused by limited ankle flexibility puts a dangerous amount of tension on the posterior muscles of the thigh as well as the adductors magnus and gracilis, increasing the risk of muscle tears. Furthermore, the lowering of the femurs below horizontal forces the lower back to round, increasing the danger of vertebral injuries.

This squat demands enormous concentration in order to maintain the correct position. Because of its risks, the use of heavier weights is limited.


Notice that stiff ankles also limit the flexion of the legs even though the thighs arrive at horizontal.

INITIAL POSITION OF THE DEADLIFT DEPENDING ON THE DIFFERENT MORPHOLOGIES



Finally, the greater cantilever than in the classic squat forces the quadriceps to provide a greater force to extend the legs.

INFLUENCE OF ANKLE FLEXIBILITY ON THE EXECUTION OF THE SQUAT



Comment: Although the femurs of figures 1 and 2 are horizontal, flexion of the thighs of figure 2 is of lesser amplitude than the flexion of the thighs of figure 1.

Improving Your Squat Position

To limit the stress on the low back and hamstring muscles, it is possible, as the powerlifters do, to lower the bar to the level of the posterior deltoids. This technique allows you to reduce the cantilever and increase the power of the lift from the back, which allows you to take on more weight.

Using a block under your heels or weightlifting or powerlifting shoes with rigid raised heels reduces the cantilever by limiting the backward movement of the gluteals as the knees move forward, allowing for greater amplitude of flexion of the knees. This technique enables you to feel the work of the quadriceps muscles while limiting the tilt of the torso and the work of the gluteus maximus muscles and the erector spinae.

The combination of bar placement and raised heels allows you to take on significantly heavier weight. It is recommended for people with long limbs or rigid ankles.

Squat With the Bar in Front to Target the Quadriceps


By limiting the tilt of the torso, the squat with the bar in front reduces the work of the low back and limits the stress on the hamstring muscles and adductor magnus.

However, by increasing the cantilever, the anterior position of the bar forces the quadriceps to work harder to extend the thigh over the lower leg.

Thus, this is the premier squat for the thighs. However, it always works with much lower weight than the classic squat. For greater stability, always execute it with raised heels.

Unfortunately, this squat is difficult for people with long limbs. A more inclined torso makes it more difficult to hold the bar, which you might drop by falling forward.


COMPARISON BETWEEN A LOW-BAR SQUAT WITH HEELS RAISED AND A CLASSIC SQUAT (WITH A LONG-LIMBED BODY)



In powerlifting, the wide-leg squat limits flexion of the torso and reduces flexion of the hips.




COMPARISON BETWEEN A SQUAT WITH THE BAR IN FRONT AND A CLASSIC SQUAT



A big belly acts as a balloon pinched between the thigh and the chest and prevents tilting of the back.





Spreading the Legs to Raise the Torso

With the squat, in order to limit excessive and dangerous tilting of the torso, you can position the legs wider apart by turning the toes out. Certain powerlifters take this technique to the extreme by placing the legs in almost a full split (which also allows them to limit their leg flexion).

The squat with widespread legs requires good flexibility of the adductor muscles of the leg and an adequate structure of the coxofemoral articulation. Therefore, not everyone can perform this type of squat.

The Advantage of a Big Belly

A big belly presses against the thighs and helps limit flexion of the hips and rounding of the back. It also protects the small of the back and limits the risk of disc herniation. This physical characteristic is common among many powerlifters and heavyweight lifters.

The Different Kinds of Knees


In weightlifting, it is important to take into account individual variations of morphology, especially in the knees.

Although bow-leggedness (genu varum) does not present more risk than normal legs, knock-kneed legs (genu valgum) or hyperextended knees (genu recurvatum) may even be a contraindication to weightlifting with heavy weights.

Genu valgum is generally found in these people:

1. Those who were overweight during their youth. During that time, the bones of the legs are not yet fully developed and still malleable, so they became deformed and develop the X shape.
2. Women, who tend to have wider hips due to their reproductive function. This can increase the angle of the femurs.

If the genu valgum is too great, the articulation is overused. The medial collateral ligament is overstretched and the lateral meniscus along with the articular surfaces covered with cartilage of the lateral condyle of the femur and the lateral external tuberosity of the tibia are subjected to excessive friction, which can lead to overuse pathologies.



TYPE A: Long legs, short torso

TYPE B: Short legs, long torso

Comment: It is important to take into account the relationship between the length of the torso and the length of the leg.

Type A: People whose legs are proportionately longer than the torso will find it more difficult to perform a squat correctly without leaning too far forward with the chest.

However, limiting the cantilever helps a person with a short torso execute the good morning, the classic deadlift, and the stiff-legged deadlift.

Type B: People whose torsos are proportionately longer than the legs will find it easier to perform the squat safely without excessively leaning forward. It is therefore not surprising that the greatest powerlifting champions specializing in the squat have this type of morphology.


Genu recurvatum is mainly found in people who are very supple (referred to as hypermobile), especially women, in whom this frequent ligamentous and muscular hyperlaxity is directly related to the reproductive function.

Rarely pathological, the recurvatum knee can involve complications such as a pinched meniscus, which occurs when the knee moves rapidly into hyperextension and the menisci do not have the time to slide, or during exercises with heavy weights, which force hyperextension of the thigh.

For this reason, people who have pathological recurvatum should never completely lock the knees at the end of extension during the squat or leg press.



DISC HERNIATION




**SCIATIC NERVE
AND POSTERIOR CUTANEOUS NERVE OF THIGH**



INCORRECT POSITION



To prevent injury at the lumbar level, never round the back when executing a deadlift or squat.



Vertebral flexion using weights can cause a herniated disc, usually in the lumbar vertebrae. These herniations occur most frequently with the squat and deadlift, and most often are the result of incorrect back position because of bad technique.

Disc herniation is a relatively frequent injury in weightlifting, most often caused by incorrect back position during the squat, deadlift, or bent row.

When executing these exercises, the main thing to avoid is rounding the back (vertebral flexion), which expands the back of the disc and pinches the front of it.

If the intervertebral disc is cracked or aging, the gelatinous liquid of the nucleus pulposus migrates backward and can compress on the spinal cord or the roots of the spinal nerves. Symptoms depend on the type of lesion, the amount of nucleus pulposus pushed out, and the surface that is compressed. The nucleus pulposus can bulge or, worse, explode through the annulus fibrosus, which surrounds it, and sometimes tear the posterior ligament that connects the vertebrae to each other. Compression of the neural elements caused by the tearing of the annulus fibrosus is particularly painful and incapacitating.

In weight training, herniations usually occur at the lumbar level and most frequently between the third and fourth or between the fourth and fifth lumbar vertebrae. The pain is dull and deep, sometimes accompanied by swelling and tingling. The pain is located in the middle of the back or more often to one side, radiating to the gluteal muscles, pelvis, and pubis and down the leg following the path of the sciatic nerve (hence the name sciatica to define this type of pain). Generally, disc herniations are spontaneously reabsorbed, and the pain eventually disappears. But in some cases, the bulge in the disc does not disappear and continues to press painfully against the nerves, or a detached piece of intervertebral cartilage compresses the neural elements. In both these cases, a surgeon can remove the part that is pressing against the nerves.


To prevent disc herniation, use proper form when performing risky exercises such as the squat, deadlift, good morning, and bent row.

Attention: No matter what the exercise, as soon as heavy weights are involved, it is essential to create a *block*.

1. Expanding the chest and holding a deep breath fills the lungs, which supports the rib cage and prevents the chest from collapsing forward.
2. Contracting the abdominal muscle group supports the core and increases the intra-abdominal pressure, which prevents the torso from collapsing forward.
3. Finally, arching the low back by contracting the lumbar muscles positions the spinal column in extension. These three actions together are referred to as *blocking*, which keeps you from rounding the back (vertebral flexion). A rounded back when lifting heavy weights can cause a herniated disc.

Comment: After a heavy workout, stretch the back by hanging from a chinning bar and focusing on relaxing the body. This allows the muscles to relax and rebalance the pressures inside the intervertebral discs.

EFFECT OF AGE ON DISC HERNIATION



LUMBAR VERTEBRAL SEGMENT (CUT)

- [1] Young vertebral segment:**
The intervertebral disc is still healthy.



- [2] Older vertebral segment:**
With age, the annulus fibrosus begins to develop fissures, and the viscous gel of the nucleus pulposus begins to dehydrate. The intervertebral disc collapses and the vertebral segments lose their mobility.

From the age of 30, the intervertebral discs degenerate, and the annulus fibrosus can crack as the nucleus pulposus begins to dehydrate. The discs of older athletes are more rigid and less elastic, and the mobility of the spine is limited. On the other hand, as the viscous gel of the nucleus pulposus gradually dehydrates, it becomes less likely that it will be displaced and compress against the nerve.


In comparison, disc herniation in a young person involves the movement of a greater amount of the gelatinous fluid of the nucleus pulposus, causing more compression, pain, and incapacity of the neural elements. Disc herniation therefore occurs more frequently with young athletes.

INCLINE LEG PRESSES

6



Attention: For some people, using the press with heavy loads can provoke a movement of the sacroiliac hinge. This can induce very painful muscle spasms.



INITIAL POSITION

Position the back properly against the backrest on the machine, with the feet slightly apart:

- Inhale and release the safety bars, then bend the knees completely so that the thighs touch or nearly touch the torso.
- Return to the initial position, exhaling at the end of the movement.

Placing the feet low against the foot plate isolates the quadriceps; placing the feet higher on the foot plate calls on the gluteal muscles and the hamstrings. Placing the feet wider apart focuses effort on the adductors.

Comment: People with back pain who are unable to perform squats can do this exercise. However, they must never lift the back off the back pad.

Feet high on the foot plate



Strong use of the gluteal muscles and the hamstrings

Feet low on the foot plate



Strong use of the quadriceps

Feet apart



Strong use of the adductors


Feet close together




Strong use of the quadriceps

7

HACK SQUATS



THE MOVEMENT



Attention: To protect the back, contract the abdominal core, which eliminates lateral movement of the pelvis or cervical spine.

Stand with the legs straight and feet slightly apart, back against the back pad, and shoulders positioned under the shoulder pads. (Hack refers to a "yoke." The pads are reminiscent of the collar placed around the neck of draft animals.):

- Inhale and release the safety catch.
- Bend the knees, then return to the initial position.
- Exhale at the end of the exercise.

This movement focuses the effort on the quadriceps. The more the feet are forward, the more the gluteal muscles will be used.

BIPEDAL ADAPTATION



Human



Chimpanzee


In the chimpanzee, our closest relative, the well-developed torso is paired with an underdeveloped gluteus maximus, which makes raising the trunk and standing erect difficult and causes an awkward two-footed gait.

The human is the only primate who has completely adapted to walking upright on two legs. Besides the well-developed gluteus maximus, the entire human structure has adapted to walking on two legs. For example, the torso is relatively small, which makes holding it erect easier, and, unlike the gorilla or chimpanzee, humans can lock the knee joint when it is extended, which makes standing much less tiring.

BOX SQUATS

8

THE MOVEMENT



1 Initial position.

2 Sit on the bench. Keep thighs relaxed and back slightly arched and always tilted forward.

The box squat is mainly used by powerlifters with the goal of strengthening the intensity of the squat.

This technique involves performing a squat by sitting on a bench for one or two seconds before standing. In a classic squat, the tension accumulated in the muscles during the negative (lowering) phase, as in stretching an elastic, is released during the positive (standing up) phase. Sitting down on a bench during the box squat relaxes the muscles of the thighs so that they cannot use the accumulated energy from the descent for the ascent.

For this reason, with the same amount of weight, the effort of the quadriceps is more intense in the classic box squat. Thus, it is a very good movement for focusing the work onto the thighs.

This movement can be included in a program for long-legged athletes who have difficulty feeling the quadriceps work during a squat. Furthermore, initiating a squat from a sitting position helps develop the automatic pushing reflex in the classic squat, making the positive standing phase quicker and more powerful.




Attention: Although the box squat is an excellent movement, you must perform the box squat very carefully, always controlling the lowering in order to sit gently onto the bench. If the lowering phase is too rapid and the gluteals slam down on the bench, the shock and the excessive pressure on the vertebral articulations can cause serious trauma.



Comment: Special benches are adjustable in height and adaptable to individual morphologies. Their cushioned seats minimize the shock of the lowering phase and limit the risk of vertebral injury by compression. To perform the movement properly, always keep the back slightly tilted. If the back is too straight when you're standing up from the bench, it will be impossible to perform the exercise.

9

LEG EXTENSIONS



THE MOVEMENT




Sit at the machine and grasp the handles or the seat to hold the torso immobile. Bend the knees and place the ankles under the ankle pads.


- Inhale and raise the legs to horizontal.
- Exhale at the end of the exercise.

This is the best exercise for isolating the quadriceps. The greater the angle of the backrest, the farther toward the back the pelvis rotates. This exercise stretches the rectus femoris (the midline biarticular portion of the quadriceps), which makes the work on it more intense while extending the legs.

This movement is recommended for beginners so that they can develop enough strength to move on to more technically demanding exercises.





INSERTIONS OF THE QUADRICEPS FEMORIS MUSCLE ONTO THE FEMUR



QUADRICEPS FEMORIS MUSCLE

STRETCHING THE QUADRICEPS



Stand on one leg:

- Grasp the foot or ankle of the other leg.
- Try to pull the heel up to the buttock.



This exercise stretches the quadriceps and, to a lesser degree, the tensor fasciae latae, as well as deeper in the iliopsoas.

To really feel the stretch on the rectus femoris (the biarticular portion of the quadriceps), pull the thigh back as far as possible. Its extension is naturally limited by the tension of the iliofemoral ligament.

Comment: For better balance, you can use the other arm to support you against a wall or stable object.

10 LYING LEG CURLS

VARIATION WITH A DUMBBELL HELD BETWEEN THE ANKLES




Lie facedown on the machine. Grasp the handles, extend both legs, and position the ankles under the ankle pads:

- Inhale and bend both legs at the same time, trying to touch the gluteal muscles with the heels.
- Exhale at the end of the effort.
- Return to the initial position with a controlled movement.


This exercise works the hamstring group and gastrocnemius and deeper, the popliteus muscle. In theory, during flexion, it is possible to target the semitendinosus and semimembranosus by internally rotating the feet, or to target the long and short heads of the biceps femoris by externally rotating the feet. But in practice, this proves to be difficult, and only emphasis on the hamstrings and the gastrocnemius can be easily achieved:

- Point the toes (plantar flexion) to feel the effort in the hamstrings.
- Flex the feet (dorsiflexion) to feel the effort in the gastrocnemius.

Variation: This exercise may be performed by alternating the legs.






HAMSTRING MUSCLES



STANDING LEG CURLS


11



Contracting the hamstrings and the lateral and medial heads of the gastrocnemius flexes the knee joint.

Hamstrings

Gastrocnemius



THE MOVEMENT


Stand with the torso resting against the pad, the thigh positioned against the knee pad, the knee straight, and the back of the ankle resting against the ankle roll:

- Inhale and bend at the knee.
- Exhale at the end of the movement.


This exercise uses the hamstring group (semitendinosus and semimembranosus and the long and short heads of the biceps femoris) and, to a lesser extent, the gastrocnemius. To engage the gastrocnemius more, simply bend at the ankle when bending at the knee. To decrease its participation, which is often the goal, simply point the toes.

SHORT HEAD OF THE BICEPS FEMORIS

Of all the flexor muscles, only the short head of the biceps femoris works across just one joint: It bends the knee.



12 SEATED LEG CURLS




Sit at the machine with the legs extended, ankles resting on the ankle pad, and thighs positioned between the thigh pad and the seat. Grasp the handles:

- Inhale and bend the knees.
- Exhale at the end of the movement.


This exercise uses the hamstring muscles and, deeper, the popliteus. To a lesser extent, it works the gastrocnemius.

Comment: The seated position (that is, with the pelvis in a forward tilt) stretches the semimembranosus, the semitendinosus, and the long head of the biceps femoris, allowing you to really target the work onto this muscle group.

POPLITEUS MUSCLE




The popliteus, located on the posterior side of the leg at the knee joint, works with the hamstrings and gastrocnemius to bend the leg.



Variations

- Perform the exercise with the feet in dorsiflexion (feet flexed) to increase the work of the gastrocnemius muscles.
- Perform the exercise with the feet in plantar flexion (toes pointed) to focus the effort on the hamstrings.




MUSCLES USED



HAMSTRING MUSCLE TEARS

ACTION OF THE HAMSTRING MUSCLES DURING THE SQUAT




2 Straightening the pelvis erects the torso.

1 The hamstrings contract to straighten the pelvis.

While performing the squat, the hamstring muscles contract to straighten the pelvis, which at the same time prevents the torso from tilting too far forward (as long as the pelvis is aligned with the torso by contracting the abdominal and lumbar muscles).

HAMSTRING MUSCLES



In bodybuilding, hamstring tears occur frequently. This injury occurs most often during the squat when the torso is too far forward. The hamstring muscle group, with the exception of the short head of biceps femoris, is in an extremely stretched position and contracts forcefully to straighten the pelvis. This can lead to tearing, most often at the high or middle portion of the muscle group.

Hamstring tears can also occur when working at a leg curl machine using heavy weights. This most often occurs at the beginning of the movement when the legs are extended and the muscles are stretched.

Although in general, the tears in hamstring muscle fibers are not extensive and not serious (it is rare to see a significant tear in the muscle or its tendinous insertion), they are always painful and prone to complications.

In fact, fibrous scarring frequently occurs after a tear in this muscle group, which creates friction that is especially painful and incapacitating during sport activity. Furthermore, this inelastic scar tissue is liable to tear during intense effort.

Preventing Hamstring Tearing


To prevent muscle tears, perform either a specific stretching workout or incorporate hamstring stretches during a lifting workout between sets of squats and deadlifts and exercises for the back of the thigh.

Certain weightlifting exercises, such as good mornings or stiff-legged deadlifts, can be considered the best hamstring protectors because of their combined action of muscle strengthening and stretching.

After Hamstring Tearing

To prevent the formation of fibrous scar tissue in the hamstrings, it is essential to reeducate the muscles as soon as possible. A week after a tear, you must perform gentle stretches for the back of the thighs. The goal is to stretch the injured muscles and especially to soften the scar so that it doesn't tear when you resume training.

RETRACTION OF THE HAMSTRINGS



In today's modern world, sitting for long periods during the day can lead to retraction of the hamstring muscles in certain people.


This retraction of the muscles on the back of the thigh tips the pelvis back and reduces the normal curvature of the spine.

This causes the person to adopt poor posture with the pelvis tucked under and the back rounded, which over time can lead to vertebral injuries. To limit this relatively frequently occurring retraction of the hamstrings, stretching movements such as an easy good morning with straight legs and the stiff-legged deadlift are recommended. Hamstring stretches after a hamstring workout are also recommended.

Retraction of the hamstrings causes the pelvis to tip back (posterior rotation), which reduces the lumbar curve and sets the stage for vertebral injuries.

Comment: A massage therapist can also treat fibrous scars by using massage or mechanical techniques aimed at softening the lesion.

13 GOOD MORNINGS



- 1 Straightening the legs while tilting forward lengthens the hamstrings, helping to better feel the contraction while straightening the trunk.
- 2 Bending the knees while tilting forward allows the hamstrings to relax, which facilitates hip flexion.




Stand with the feet slightly apart and the bar resting on the trapezius or a little lower on the posterior deltoid:

- Inhale and bend the torso forward, keeping the back straight. The axis of rotation should pass through the coxofemoral joints.
- Return to the initial position and exhale.


To make the exercise easier, bend slightly at the knees.

This movement, which works the gluteus maximus and the spinal group, is especially noteworthy for the action on the hamstrings (except the short head of the biceps femoris, which only flexes the knee). Besides knee flexion, the main function of the



TWO WAYS TO PERFORM GOOD MORNINGS

- 1 Bent knees
- 2 Straight knees



hamstrings is to tip the pelvis back (posterior rotation) and straighten the torso when the pelvis is locked to the torso through isometric contraction of the abdominal core and the lumbosacral muscle group.

To better feel the work of the hamstrings, don't work with heavy weights. In the negative phase, the good morning is excellent for stretching the back of the thighs. Worked regularly, it helps prevent injury when executing a heavy squat.

STRETCHING THE HAMSTRINGS



Stand on one leg, slightly bending at the knee; the other leg is extended with the foot in dorsiflexion:


- Place the hands on the thighs and slightly arch the back. Slowly bend the torso forward, concentrating on the feeling of stretching at the back of the thigh. The tilt occurs at the pelvis.
- Maintain this position for 20 seconds. Return slowly to the initial position and change sides.

This exercise mainly stretches the hamstring muscle group and adductor magnus as well as the gastrocnemius, soleus, and, to a lesser degree, gluteus maximus.

Variation: Stand on one leg with the other leg extended and resting on a bench with the foot in dorsiflexion:

- Place the hands on the thigh of the leg that is extended. Slightly arch the back and slowly tilt the torso forward, focusing on the feeling of stretching at the back of the thigh. The tilt occurs at the pelvis.
- Maintain the position for 20 seconds. Return slowly to the initial position and change sides.

To better focus on the stretching of the hamstrings, relax the calf muscles by putting the front foot into plantar flexion.



Comment: To avoid injury in the squat and the deadlift, practice this stretch at the beginning of a session by incorporating it among the first series.


Attention: In weightlifting, the main function of stretching movements is to equalize the muscle fiber tension inside the muscle and limit the risk of injury. With heavy weights, if the muscle fiber tension is not homogeneous, the tightest fibers risk tearing.

For this reason, at the beginning of the training session during the first warm-up series, perform a few stretching exercises specific to the muscle groups that will be worked.

Always perform the stretching movements gently and in moderation to protect the articulations and to avoid excessive stretching of the ligaments. Excessive and aggressive stretching risks destabilizing the articulation and generating pathological inflammation.





VARIAION WITH A BENCH



When bending forward at the pelvis (pelvic tilt), the hamstring muscles are stretched.

14 CABLE ADDUCTIONS




Stand on one leg, with the other leg in the ankle cuff and the hand of the supporting leg holding onto the machine:

- Pull the cable across the support leg.


This exercise works the adductor group (pectineus; adductors brevis, longus, and magnus; and gracilis). To develop definition of the inside of the thighs, perform sets of high repetitions.

ADDUCTOR MUSCLES OF THE THIGHS



MACHINE ADDUCTIONS

15



Sit at the machine with the legs spread apart:


- Contract the thighs to bring the legs together.
- Return to the initial position with a controlled movement.

This exercise works the adductor muscle group (pectineus; adductors minimus, magnus, brevis, and longus; and gracilis) and allows you to use heavier weights than you can with the cable adductions, but with a decreased range of motion.

Best results are achieved in long sets until you feel a burn.


Comment: Perform this exercise to strengthen the adductors. This muscle group is often injured during intense exertion.

Therefore, increase the weights gradually and perform adductor muscle stretches at the end of the workout.




THE MOVEMENT

ADDUCTORS MINIMUS AND MAGNUS (POSTERIOR VIEW)



16

FLEXION AND EXTENSION OF THE FEET




Stand on a step with one hand holding a wall or railing for stability:

- Slowly dorsiflex your feet (lower your heels) to get a good stretch in the calves.
 - Then plantar flex (rise up on your toes) while keeping the knees extended or slightly bent.
- Perform this exercise slowly in a long series until you feel a burn.

The combined action of muscle contraction and stretching makes this movement ideal as a warm-up at the beginning of a training session for the calves with the purpose of avoiding injury, or at the end of the session in order to really feel the muscle contraction.



This exercise mainly works the triceps surae (made up of the two gastrocnemius and the soleus) as well as the flexor hallucis longus, tibialis posterior, and flexor digitorum longus (these last three muscles are located deeper).

Comment: This movement is also excellent for stretching the muscles of the plantar surface of the foot, such as the flexor digitorum brevis and the quadratus plantae, and for making the plantar aponeurosis more supple.



STANDING CALF RAISES

17

GASTROCNEMIUS MUSCLE
MEDIAL HEADGASTROCNEMIUS MUSCLE
LATERAL HEAD



Triceps surae

- Gastrocnemius, lateral head
- Gastrocnemius, medial head
- Soleus

Tibia, medial malleolus
Fibula, lateral malleolus
Calcaneal tuberosity



VARIATION
STANDING BARBELL CALF RAISES



1 Begin 2 End

THE MOVEMENT



VARIATION

Performing the exercise at an incline machine works the calves without overloading the back.




Stand at the machine with a straight back, shoulders under the pads, and the balls of the feet on the foot plate, with the calves relaxed and the heels hanging down:

- Rise up by extending (plantar flex) the feet, keeping the knees straight.

This exercise uses the triceps surae (made up of the soleus and the lateral and medial heads of the gastrocnemius). Move the feet through the complete range of flexion with each repetition in order to stretch the muscles properly. In theory, it is possible to isolate the medial gastrocnemius by pointing the toes out and to isolate the lateral gastrocnemius by pointing the toes in. But in practice, this is difficult to achieve. Only separating the work of the soleus and gastrocnemius is easy to achieve. This is done by flexing the knees to relax the gastrocnemius and to put more effort on the soleus.

Variations: Perform the exercise at a frame with a wedge under the feet or with a free bar without the wedge for more balance; however, this reduces the amplitude of movement.

Comment: The triceps surae is an extremely powerful, tough muscle group that alone raises the entire weight of the body thousands of times in a day when we walk. Don't hesitate to work it with heavy weights.




TRICEPS SURAE ACTION

18 ONE-LEG TOE RAISES




INITIAL POSITION




Stand with the toes of one foot on the foot plate and hold a dumbbell in one hand and use the other hand for support and balance:

- Rise up on the toes (plantar flexion), keeping the knee joint straight or slightly flexed.
- Return to the initial position.

This exercise contracts the triceps surae. Completely flex the foot with each repetition in order to stretch the triceps surae properly. Optimal results are obtained through long sets until you feel a burn.




TWO TYPES OF CALVES




Comment: Some people have an unusual triceps surae that does not grow larger with training. These people can develop strength only. Long gastrocnemius and soleus muscles develop easily. Conversely, short calves resist developing bulk.

TRICEPS SURAE MUSCLES




DONKEY CALF RAISES

19



TRICEPS SURAE INSERTION




Stand with the legs relaxed, toes on the foot plate and heels hanging down, legs extended, torso leaning forward, forearms resting on the front support, and the padded plate of the machine resting on the back of the hips:

- Rise up by extending the ankles and pointing the toes (plantar flexion).



This exercise focuses on the triceps surae, especially the gastrocnemius.

Variation: If there's no machine for this exercise, put a block under the feet, bend forward, and rest the forearms on a support, and perform this exercise while someone straddles your hips.



TRICEPS SURAE ACTION



20 SEATED MACHINE CALF RAISES



Quadriceps
– Rectus femoris
– Vastus lateralis
– Vastus intermedius
Tensor fasciae latae
Fascia lata, iliotibial tract



Sit at a machine with the knees positioned under the pads, the toes on the foot bar, and the ankles relaxed:


- Extend the feet and point the toes.

This exercise isolates the soleus, whose name is derived from its resemblance to the flat fish, the sole.

(This muscle inserts at the top at the tibia and fibula under the knee joint and attaches at the bottom to the calcaneus by the Achilles tendon. Its purpose is to extend the feet at the ankles.)

Bending at the knees relaxes the gastrocnemius, which attaches at the top above the knee joint and at the bottom onto the Achilles tendon, and reduces its contribution to ankle extension.

Variation: You can also perform this exercise by sitting on a bench with a wedge under the feet and a barbell resting on the thighs. Wrap the bar for comfort.





SEATED BARBELL CALF RAISES

21



INITIAL POSITION



Sit on a bench with a block under the toes and the barbell resting on the thighs:


- Extend (plantar flex) the ankle.

This exercise mainly uses the soleus. This muscle, which is part of the triceps surae group, inserts at the top below the knee joint on the tibia and fibula. At the bottom, it attaches to the calcaneus via the Achilles tendon. Its function is to extend the feet at the ankles. Unlike the calf raises, which allow you to work with heavy weights, this exercise does not allow heavy weights because of the awkwardness of the bar position. For best results, work in sets of 15 to 20 repetitions.



Attention: Cushion the bar on the thighs with a rubber pad or a folded towel to reduce pain.

Variation: You can also perform this exercise without additional weights while sitting on a chair or bench. In this case, work in very long sets until you feel a burn.




When the knees are bent, the gastrocnemius muscle, which attaches above the knee, is relaxed. In this position, it weakly assists ankle extension because most of the work is done by the soleus.

Conversely, when the knee is straight, the gastrocnemius is stretched. In this position, it actively participates in ankle extension and completes the action of the soleus.



STRETCHING THE CALF



Stand with your hands on your hips, with one leg forward in a lunge stance, the other leg extended behind, and your feet in line with the knees:

- Bend the forward knee by advancing with your pelvis, being careful to always keep the back leg extended with the heel on the ground.
- Hold this position until you feel the stretch on the posterior leg.


This exercise mainly solicits the triceps surae, which is made up of the gastrocnemius and the soleus; the flexor digitorum muscles; and the tibialis posterior located deeper beneath the triceps surae. To a lesser degree it solicits the peroneus longus and brevis.



SHORT CALF, LONG CALF

There are great individual differences in the shape of calves. A large part of these morphological variations developed during the first human migrations as evolutionary adaptation to climate. Thus Black Africans commonly have legs that are proportionately longer, a dropped plantar arch (flat feet), and a relatively long calcaneus. This osseous configuration of the leg and foot creates an excellent lever at the ankle with a minimum of triceps surae muscle (the muscle of the calf is short, slender, and high with a long tendon), which produces powerful extension of the foot while walking. On the other hand, Nordic people frequently have legs that are proportionately shorter with a very prominent plantar arch and a short, close calcaneus. This osseous conformation of the leg and foot with a shorter lever requires a voluminous and not very economical triceps surae that reaches far down for extension while walking. This type of long, bulky calf is in fact adapted for cold climates. It conserves the heat of the body, offering a minimal exterior surface and limiting the thermal exchanges and incapacitating, if not fatal, chilling during periods of intense cold. Although often considered more attractive, the long and bulky calf is less well adapted for running and more vulnerable to muscle tearing. Thus, this type of calf needs a more careful warm-up and requires stretching movements before and after intense training sessions.

THE TWO KINDS OF CALVES



6

BUTTOCKS

Gluteal Muscles, a Human Characteristic


Although some of the larger primates occasionally walk, humans are the only primates and one of the few mammals that have completely adapted to two-legged locomotion. One of the structural features directly related to this way of getting around is the significant development of the gluteus maximus muscle, which has become the biggest and most powerful muscle in the human body.

The development of the gluteal muscles is truly a human characteristic. In comparison, the gluteal muscles in quadrupeds are proportionately underdeveloped, and the hindquarters of the horse, which some consider as typical for animals, are in fact made up of the hamstrings (the back of the thigh in humans).

In humans, the gluteus maximus, which extends the hip, does not play an important role in walking. Instead, the hamstrings play the major role in straightening the pelvis (hip extension) with each stride. Just put your hand on the buttocks while walking, and you can feel that they do not contract much.

However, as soon as the effort becomes significant, such as when walking uphill, walking quickly, or running, the gluteal action is called into play to extend the hip and erect the torso.


These biomechanical points help explain why in exercises for the gluteal muscles and the hamstrings, such as good mornings (see page 144) and stiff-legged deadlifts (see page 102), either the gluteal muscles or the hamstrings are isolated depending on the amount of weight involved.



| | |
|--|------------|
| 1 BARBELL LUNGES | 156 |
| 2 DUMBBELL LUNGES | 157 |
| + KNEE INSTABILITY | 158 |
| 3 CABLE BACK KICKS | 159 |
| 4 MACHINE HIP EXTENSIONS | 160 |
| 5 FLOOR HIP EXTENSIONS | 161 |
| 6 BRIDGING | 162 |
| + STRETCHING THE GLUTEUS MAXIMUS AND HAMSTRINGS | 163 |
| 7 CABLE HIP ABDUCTIONS | 164 |
| + INDIVIDUAL VARIATIONS IN HIP MOBILITY | 165 |
| 8 STANDING MACHINE HIP ABDUCTIONS | 166 |
| 9 LYING HIP ABDUCTIONS | 167 |
| 10 SEATED MACHINE HIP ABDUCTIONS | 168 |
| + STRETCHING THE GLUTEALS | 169 |

1

BARBELL LUNGES




Stand with the legs slightly apart and the bar behind the neck resting on the trapezius muscles:

- Inhale and take a big step forward, keeping the trunk as straight as possible.
- Lunge until the front thigh is horizontal to the floor or slightly less.
- Exhale and return to the initial position.

This exercise, which works the gluteus maximus intensely, can be performed two different ways: either by taking a small step (which isolates the quadriceps) or taking a big step (which isolates the hamstrings and gluteus maximus and stretches the rectus femoris and iliopsoas of the back leg).

Comment: Because the front leg must support almost all the weight in the lunge position and the exercise demands a good sense of balance, begin with very light weights.




1 Execution with a small step predominantly works the quadriceps.

2 Execution with a big step predominantly works the gluteus maximus.

DUMBBELL LUNGES

2



Stand with the legs slightly apart and hold a dumbbell in each hand:

- Inhale and take a big step forward, keeping the torso as straight as possible.
- When the forward thigh reaches horizontal or slightly below, use tonic extension to return to the initial position.
- Exhale at the end of the movement.


This exercise mainly works the gluteus maximus and quadriceps.

The bigger the step, the more the gluteus maximus of the forward leg is used and the iliopsoas and rectus femoris of the back leg are stretched.

A smaller step isolates the quadriceps of the forward leg.

You can perform a complete set on one side and then the other or work the legs alternately during the same set.

Comment: Because all of the weight is supported by the front leg in the lunge position and the exercise requires a good sense of balance, work with light weights to protect the knee.





KNEE INSTABILITY


When the knee is extended, the medial and lateral collateral ligaments are stretched and prevent rotation of the joint. When you are standing, the knee locks in extension, and there is no need for muscle tension to stabilize the joint.

When the knee is bent, the medial and lateral collateral ligaments are relaxed. In this position, muscle tension provides the stability.

When the knee flexes and rotates, the meniscus travels forward. Then, if extension is not controlled, the meniscus may not return to its normal position fast enough and becomes pinched between the condyles, which can tear the meniscus. If a piece of the meniscus is severed when it is pinched, surgery may be necessary to remove it.


With asymmetrical exercises such as the lunge (see pages 156 and 157), control the speed and the form of the movement to protect the knee.

1 KNEE IN EXTENSION




When the knee is flexed, the lateral ligaments are relaxed. Rotation of the joint is possible.

2 KNEE IN FLEXION




MENISCI




One of the main functions of the meniscus is to disperse pressure in the knee joint by increasing the supporting surface area for the femur on the tibia, avoiding premature wear on the articular surfaces.

MENISCI AND KNEE LIGAMENTS




CABLE BACK KICKS



3



The deep gluteus maximus attaches to the linea aspera of the femur.



Hip extension is limited by the iliofemoral (Bertin's) ligament.



Stand on one leg facing the machine, the other leg attached to the ankle strap of the low pulley, and the pelvis tilted forward. Grasp the handle:


- Extend the hip and pull the leg back.
- Hip extension is limited by the tension of the iliofemoral (Bertin's) ligament.

This exercise mainly works the gluteus maximus and, to a lesser extent, the hamstrings (except the short head of the biceps femoris).




It helps develop the profile of the hips while firming the gluteal region.

4

MACHINE HIP EXTENSIONS



Hip extension is limited by the tension of the iliofemoral ligament, also referred to as Bertin's ligament, which is a thickening of the articular capsule.






Stand on one leg with the other slightly forward and position the pad against the calf halfway between the knee and ankle. Lean the torso forward slightly and grasp the handles:

- Inhale and push the thigh back until the hip is hyperextended.
- Maintain the position with an isometric contraction for a couple of seconds.
- Exhale at the end of the extension.
- Return to the initial position.

This exercise mainly works the gluteus maximus and, to a lesser extent, the semitendinosus, semimembranosus, and the long head of the biceps femoris.

FLOOR HIP EXTENSIONS

5



THE MOVEMENT

Kneel on one leg and bring the other knee to the chest while leaning on the elbows or on the hands with the arms extended:

- Extend the bent leg back with complete hip extension.

With the leg extended, this exercise uses the hamstrings and gluteus maximus. With the knee bent, only the gluteus maximus is used, and less intensely.


This exercise can be performed with higher or lower amplitude during the last part of the extension. You can maintain an isometric contraction for a couple of seconds at the end of the movement.

To increase the intensity, use ankle weights.



Its ease of execution and its effectiveness have made this exercise popular, and it is frequently used in group classes.

1 Begin

2 End



VARIATION ON A BENCH




VARIATION WITH BENT KNEE

6

BRIDGING

THE MOVEMENT



Lie on the back with hands flat on the ground, arms alongside the body, and knees bent:

- Inhale and lift the buttocks off the ground, pushing down through the feet.
- Maintain the position for a couple of seconds and lower the pelvis without touching the buttocks to the ground.
- Exhale and begin again.


This exercise mainly works the hamstrings and gluteus maximus.

Perform this exercise in long sets, making sure to contract the muscles at the top of the lift, when the pelvis is off the ground.


Comment: Because it is easy and effective, bridging has become part of most group exercise classes.

VARIATION AT A BENCH

1 Begin



2 End



Variation 1

To perform bridging with the feet raised, lie on the back with hands flat at the sides, arms alongside the body, thighs vertical, and feet resting on a bench:

- Inhale and raise the pelvis off the ground; maintain the position for two seconds and lower without touching the buttocks to the ground.
- Exhale and begin again.

This exercise works the gluteus maximus and especially the hamstrings. The hamstrings are used more in this exercise than when bridging from the ground. Execute this exercise slowly, and focus on the muscle contraction. Sets of 10 to 15 repetitions provide the best results.


Another variation is to perform bridging with the calves resting on the bench. This isolates the hamstrings even more intensely and also requires strong work from the gastrocnemius.

Variation 2

Limit the range of the movement by not lowering the pelvis as far to create a burn.

Comment: Bridging is actually extending the hips.


STRETCHING THE GLUTEUS MAXIMUS AND HAMSTRINGS



Lie on your back with legs extended on the floor:

- Using your hands, gently bring one leg with the knee bent to your chest (to relax the hamstring muscles).
- Maintain the position, breathing slowly and trying to feel the stretching of the gluteus maximus muscle.
- Return to the initial position; change legs.




Variation: You can perform the movement by bringing the extended knee toward the chest. In this case, the stretch will be more intense on the hamstrings and less on the gluteus maximus. Note that tension on the hamstring muscles might strongly limit bending at the hip.




7

CABLE HIP ABDUCTIONS

GLUTEAL INSERTIONS AT THE OS COXA



SECTION THROUGH A FEMALE HIP DISPLAYING THE GLUTEUS MEDIUS OVERLYING THE GLUTEUS MINIMUS




Stand on one leg with a pulley attached to the other ankle:

- Raise the leg laterally as high as possible.


This exercise mainly works the gluteus medius and the deeper gluteus minimus.

Long sets until you feel a burn are most effective.

GLUTEAL "DELTOID"



GLUTEUS MINIMUS



Although located deep, the gluteus minimus contributes to the bulk of the upper buttock.




INDIVIDUAL VARIATIONS IN HIP MOBILITY

Regardless of individual muscle elasticity and ligamentous tension, it is mainly the shape of the bones of the coxofemoral joint that is responsible for hip mobility. The configuration of the bone is most important in hip abduction.

Examples


- When the neck of the femur is almost horizontal (coxa vara) and associated with a well-developed superior rim of the acetabulum covering the head of the femur, abduction movements are limited.
- When the neck of the femur is close to vertical (coxa valga) and associated with an undeveloped superior acetabular rim, abduction movements are facilitated.

Therefore, it is useless to try to raise the leg high laterally if your hip joint is not made for it.




- 1 Abduction of the hip (limited by the neck of the femur butting against the acetabulum)
- 2 Forced abduction of the hip (tilting the pelvis onto the head of the opposite femur)


VARIATIONS IN OSSEOUS HIP STRUCTURE



An almost horizontal neck of the femur is referred to as a **coxa vara**. It limits abduction movements because it butts up against the rim of the acetabulum sooner.





An almost vertical neck of the femur is referred to as **coxa valga**. It allows greater abduction movements.



ABDUCTION IS LIMITED BY THE NECK OF THE FEMUR BUTTING UP AGAINST THE RIM OF THE ACETABULUM.

8



STANDING MACHINE HIP ABDUCTIONS



Stand on one leg at the machine and place the other leg against the pad below the knee:

- Slowly raise the leg as high as possible.
- Return to the initial position.



Abduction is limited by how soon the neck of the femur butts up against the rim of the acetabulum. This exercise develops the gluteus medius. It also develops the deeper gluteus minimus, whose function is the same as that of the anterior fibers of the gluteus medius. For best results, use long sets.




GLUTEUS MEDIUS AND MINIMUS RAISE THE LEG Laterally

LYING HIP ABDUCTIONS

9



THE MOVEMENT




Lie on the side and support the head with the hand or rest the upper body on your elbow:

- Raise the leg laterally no more than 70 degrees, keeping the knee straight. This exercise works the gluteus medius and minimus. You can vary how high you raise the leg. Hold the leg at the height of the movement for a few seconds with an isometric contraction.


You can raise the leg slightly to the front or the back or raise it vertically.

To increase the intensity, use ankle weights, an elastic band, or a low pulley.

THREE WAYS TO RAISE THE LEG





ISOLATED ZONES



- 1 Leg raised vertically
- 2 Leg raised slightly behind
- 3 Leg raised slightly in front

10 SEATED MACHINE HIP ABDUCTIONS




Sit at the machine:

- Spread the legs as wide as possible.
- Return to the initial position with a controlled movement.


The more angled the backrest, the more the gluteus medius is isolated. The more vertical the backrest, the more the gluteus maximus is worked. Ideally, lean forward or back to change the angle of the torso during a set.

Example: Perform 10 repetitions with the torso resting against the backrest and 10 repetitions with the torso leaning forward.


This exercise sculpts and firms the top of the hip, which makes the waistline look narrower.



STRETCHING THE GLUTEALS



VARIATION TO ACCENTUATE STRETCHING OF THE LUMBAR REGION




Comment: This variation stretches the internal oblique, erector spinae, and splenius capitis on the side of the leg that is on the floor. It stretches the external oblique, rotators, and multifidus of the neck as well as the sternocleidomastoid on the side of the bent leg.

Sit on the floor with one leg extended and the other bent with the foot on the floor outside the extended leg. Press to the inside against the external surface of the knee of the bent leg with the opposite elbow. This exercise mainly stretches the gluteus maximus muscle and deeper in the external hip rotator group (piriformis, gemelli, quadratus femoris, and obturator internus and externus).


Variation: Rather than use the elbow to put pressure against the knee, you can clasp it with both hands.

This exercise stretches the deep, small, external rotator muscles of the hip.




7

ABDOMEN



| | |
|--|------------|
| + CORRECT POSITION FOR THE ABDOMINALS | 171 |
| 1 CRUNCHES | 172 |
| 2 SIT-UPS | 173 |
| 3 GYM LADDER SIT-UPS | 174 |
| 4 CALVES OVER BENCH SIT-UPS | 175 |
| 5 INCLINE BENCH SIT-UPS | 176 |
| 6 SUSPENDED BENCH SIT-UPS | 177 |
| 7 HIGH-PULLEY CRUNCHES | 178 |
| 8 MACHINE CRUNCHES | 179 |
| 9 INCLINE LEG RAISES | 180 |
| + STRETCHING THE UPPER BODY | 181 |
| 10 LEG RAISES | 182 |
| 11 HANGING LEG RAISES | 183 |
| 12 TRUNK ROTATIONS | 184 |
| 13 DUMBBELL SIDE BENDS | 185 |
| 14 ROMAN CHAIR SIDE BENDS | 186 |
| 15 MACHINE TRUNK ROTATIONS | 187 |
| + STRETCHING THE ABDOMINALS | 188 |


CORRECT POSITION FOR THE ABDOMINALS



ACTION OF PSOAS MAJOR ON THE LUMBAR CURVE

Aside from its role as a powerful hip flexor, the psoas muscle pulls the lumbar spine into lordosis, increasing the curve.


When performing exercises for the abdominal muscles, round the back.



CORRECT POSITION, ROUNDED BACK

INCORRECT POSITION, ARCHED BACK

As with most movements involving the abdominal core, leg raises either on the ground or at an incline bench should never be performed with an arched back.



INCORRECT POSITION, ARCHED BACK

Unlike other weightlifting movements, exercises for the abdominal core and especially those for the rectus abdominis absolutely must be worked with a rounded back (rolling up the spine).

When performing exercises that roll the spine up off the floor, as in crunches, you hold the spine differently than when performing squats, deadlifts, or other standing movements.

If during exercises with additional weights, such as squats, deadlifts, or good mornings, the vertebral column is not arched at the lumbar spine, vertical pressure combined with rounding the back pushes the nucleus pulposus of


the intervertebral disc posteriorly, which can compress the nerves and cause sciatica or a herniated disc.

On the other hand, when performing specific exercises for the abdomen, if the back is not rounded with intense contraction of the rectus abdominis and the internal and external obliques, the powerful psoas hip flexors will increase the lumbar curve, forcing the intervertebral discs forward.

This causes increased pressure at the posterior lumbar vertebral articulations, which can cause low back pain or, more seriously, articular compression or shearing.

1

CRUNCHES*





Lie on the back with hands behind the head, thighs vertical, and knees bent:

- Inhale and raise the shoulders off the ground, bringing the knees and head toward each other by crunching, which means rounding the back and rolling the spine up.
- Exhale at the end of the movement.

This exercise mainly uses the rectus abdominis.


To work the obliques more intensely, bring the right elbow to the left knee, then the left elbow to the right knee alternately with each crunch.

* Perform a crunch by rounding the back and rolling the spine up, bringing the pubis and sternum toward each other through voluntary contraction.



SIT-UPS

2



Lie on the back with knees bent, feet flat on the ground, and hands behind the head:

- Inhale and raise the torso by rounding the back.
- Exhale at the end of the movement.
- Return to the initial position without touching the ground.
- Continue until a burn develops in the abdominal muscles.


This exercise works the hip flexors as well as the obliques, but it mainly acts on the rectus abdominis.

Variations


- Having a partner hold the feet makes the exercise easier.
- Extending the arms forward makes the exercise easier for beginners.
- Working on an incline bench makes the exercise more intense.



PERFORMING THE EXERCISE WITH A PARTNER ANCHORING THE FEET



VARIATION ON AN INCLINE BENCH
The greater the angle, the greater the effort.





THE MOVEMENT

VARIATION WITH ARMS EXTENDED
To make the movement easier

Comment: Because, in general, a woman's torso is not as bulky proportionate to the legs as in men, performing sit-ups without lifting the feet off the ground is easier for women than for men.

3

GYM LADDER SIT-UPS




Lie faceup on the ground and position the feet between two bars in the ladder with the thighs vertical and hands behind the head:

- Inhale and raise the torso as high as possible, rounding the spine.
- Exhale at the end of the movement.


This exercise works the rectus abdominis and, to a lesser degree, the external oblique.

Position the feet lower on the ladder so that the pelvis can rock more and better contract the flexor muscles of the hip (iliopsoas, rectus femoris, and tensor fasciae latae) when lowering the torso.



CALVES OVER BENCH SIT-UPS

4



CORRECT POSITION




INCORRECT POSITION

PLACEMENT OF THE HANDS AND ELBOWS

To prevent pulling on the neck excessively, place the hands behind the ears rather than behind the head.

The wider the elbows, the more difficult the movement.

Conversely, the closer together and more forward the elbows, the easier the execution.



THE MOVEMENT


Lie on your back with your calves laying over a flat exercise bench. Place your hands behind your ears:

- Inhale and lift your shoulders off the floor.
- Try to touch your knees with your head.
- Exhale as you complete the movement.

This exercise focuses on the rectus abdominis, particularly above the navel. By placing your torso to move farther from the bench you increase pelvic mobility, which allows your torso to move upward by contracting the iliopsoas, tensor fasciae latae, and rectus femoris in order to flex the hips.

5

INCLINE BENCH SIT-UPS





ORIENTATION OF THE ABDOMINAL MUSCLES THAT SUPPORT AND PROTECT THE INTERNAL ORGANS

- 1 Rectus abdominis
- 2 External oblique
- 3 Internal oblique
- 4 Transversus abdominis

In quadrupeds, the muscles of the abdominal core create a hammock-like structure that passively supports the internal organs. These muscles move little during locomotion.

With the shift to bipedal locomotion in humans, the muscles of the abdominal core have grown stronger to align the pelvis with the trunk in a vertical position and to prevent the trunk from swaying too much during walking or running.

The abdominal core has developed into powerful muscles that actively contain the internal organs.



Sit on a bench with the feet positioned under the pads, hands behind the ears:

- Inhale and lower the torso less than 20 degrees.
- Raise the torso while slightly rounding the back to better focus on the rectus abdominis.
- Exhale at the end of the movement.



Perform this exercise in long sets. It works the abdominal core as well as the iliopsoas, tensor fasciae latae, and rectus femoris of the quadriceps. The latter three muscles tilt the pelvis forward.

Variation: Rotating the torso on the way up focuses some of the effort on the internal and external obliques.

Example: Rotating to the left works the right external oblique, the left internal oblique, and the right rectus abdominis more intensely. Rotations can be performed in sets of alternating sides or sets all on the same side. In either case, concentrate on feeling the muscles contract. There is no point in angling the bench excessively.



SUSPENDED BENCH SIT-UPS

6



7

HIGH-PULLEY CRUNCHES



ACTION OF THE ABDOMINAL MUSCLES




Kneel in front of the machine and hold the handle behind the neck:

- Inhale.
- Exhale and roll the spine as you lower the sternum toward the pubis.


This movement is never performed with heavy weights. Concentrate on feeling the muscles contract, mainly the rectus abdominis, in order to focus the work on the abdominal core.

MACHINE CRUNCHES

8



RECTUS ABDOMINIS MUSCLES




Sit at the machine, grasp the handles, and position the feet under the pad:

- Inhale and roll the spine, trying to bring the sternum to the pubis.
- Exhale at the end of the movement.

This exercise allows you to adjust the weight, or workload, to your ability. To avoid risk, beginners should use light weights. Well-trained athletes can use heavier weights.


9 INCLINE LEG RAISES




Lie on an incline bench and grip the bars or handles:

- Inhale and raise the legs to horizontal.
- Then raise the pelvis, rolling the spine up as if trying to bring the knees to the head.

This exercise first works the iliopsoas, tensor fasciae latae, and rectus femoris of the quadriceps when raising the legs. Then it works the abdominal core and contracts mainly the infraumbilical portion of the rectus abdominis when raising the pelvis and rolling up the spine.



Comment: This is an excellent exercise if you have trouble feeling the work on the lower abdominal muscles. Given the difficulty of the exercise, beginners should start with the bench only slightly inclined.



STRETCHING THE UPPER BODY



Stand with your legs wider apart than the pelvis and the back very straight:

- Extend the arms vertically, hands clasped together and palms facing the ceiling.
- Inhale and expand the chest, stretching the intercostal muscles and trying to push up. Keep the back and head very straight.
- Exhale slowly while relaxing, and repeat.


This is a general stretching exercise for the upper body and particularly for the intercostal muscles, rectus abdominis, latissimus dorsi, teres major, and the long portion of the triceps.

When bending to the side, the stretch becomes more intense on the external and internal obliques of the abdomen, quadratus lumborum, and the inferior and middle erector spinae muscles.


Comment: This stretch is excellent for relaxing and returning to a state of calm after an intense session of thigh presses, squats, or deadlifts when the rib cage and spine have been compressed.

It can on occasion replace or complement the stretch at a chin-up bar (see page 114) to equalize the pressures and tensions of the intervertebral articulations.



VARIATION WITH LATERAL SIDE BEND OF THE TORSO



10 LEG RAISES




THE MOVEMENT



Iliopsoas

ILIOPSOAS ACTION



Support the body by resting the elbows on the pads. Position the back against the back support:

- Inhale and raise the knees to the chest, rounding the back in order to contract the abdominal core.
- Exhale at the end of the movement.


This exercise works the hip flexors, mainly the iliopsoas, and the obliques. It intensely works the lower part of the rectus abdominis.

Variations

- To target the lower abdominal muscles, perform small flutters with the legs when rolling up the spine.
- To make the exercise more intense, extend the legs horizontally. However, this requires flexible hamstrings.
- Hold the knees to the chest for a few seconds with an isometric contraction.


HANGING LEG RAISES

11





VARIATION


Alternately raising the legs to the left and then to the right side uses the internal and external obliques more intensely.



Attention: Torso rotations are contraindicated for people who have low back problems or who have already had a herniated disc.



Hypotonic abdominal muscles create abdominal ptosis.



Hypertonic abdominal muscles.

ABDOMINAL-LUMBAR EQUILIBRIUM

Balance the work between the abdominal muscles and the erector muscles of the spine.

Hypotonicity or hypertonicity of either of these muscle groups can lead to poor posture, which over time can cause injury.

Example

Hypertonicity of the lower part of the erector muscles of the spine (lumbosacral mass) associated with hypotonicity of the abdominal muscles leads to hyperlordosis with abdominal ptosis (sagging).

If addressed in time with exercises to strengthen the abdominal core, this postural fault can sometimes be corrected.

Conversely, hypertonic abdominal muscles associated with slack erector muscles, especially in the upper part (spinalis thoracis, longissimus thoracis, iliocostalis thoracis) leads to kyphosis (rounding of the upper back) with loss of the lumbar curve. This postural fault can be corrected with exercises to strengthen the erector muscles of the spine.



Hang from a chin-up bar:

- Inhale and raise the knees as high as possible by rolling up the spine and bringing the pubis toward the sternum.
- Exhale at the end of the movement.

This exercise uses the iliopsoas, rectus femoris, and tensor fasciae latae when you raise the legs and the rectus abdominis and, to a lesser degree, the internal and external obliques when you bring the pubis toward the sternum.

Small leg flutters without lowering the knees below horizontal focus the effort on the abdominal core.

12 TRUNK ROTATIONS



Attention: Rotations of the torso are contraindicated for people who have low back problems or who have already had a disc herniation because they risk aggravating or reviving this lumbar pathology.

Stand with the legs apart. Hold a stick across the trapezius above the posterior deltoid, hands resting on the stick without pushing:


- Rotate the torso to one side and to the other, keeping the pelvis fixed with isometric contraction of the gluteal muscles.

When the right shoulder is forward, this exercise works the right external oblique and, deep in, the left internal oblique and, to a lesser degree, the rectus abdominis, quadratus lumborum, and the extensor muscles of the spine on the left side.

To increase the intensity, round the back slightly.

You can also perform the movement while seated, which helps to fix the pelvis so that you can focus the effort on the abdominal core.




Best results are obtained with sets lasting several minutes.



VARIATION SEATED ON A BENCH

DUMBBELL SIDE BENDS

13




VARIATION AT A LOW PULLEY

Stand with the legs slightly apart and one hand behind the ear. Hold a dumbbell in the other hand:

- Bend the torso to the side opposite to the dumbbell.
- Return to the initial position or beyond with passive flexion of the torso.
- Alternate sets, changing the side of the dumbbell without resting.

This exercise mainly works the obliques on the side the torso bends toward. It works the rectus abdominis, deep muscles of the back, and quadratus lumborum (back muscle that inserts on the 12th rib, the transverse processes of the lumbar vertebrae, and the iliac crest) less intensely.



QUADRATUS LUMBORUM MUSCLE

14 ROMAN CHAIR SIDE BENDS




This exercise is performed on a bench originally designed for lumbar extensions.

Lie on your side with the hip on the bench, torso in the air, hands near the ears or on the chest, and feet positioned under the pads:


- Raise the side of the body toward the ceiling.

This exercise mainly works the obliques and rectus abdominis on the side that is bending, but the opposite obliques and rectus abdominis are also used in isometric contraction to prevent the torso from lowering below horizontal.

Comment: The quadratus lumborum muscle is always used when bending the torso toward the side.




EXTERNAL OBLIQUE MUSCLE OF THE ABDOMEN



INTERNAL OBLIQUE MUSCLE OF THE ABDOMEN

MACHINE TRUNK ROTATIONS


15




Attention: Rotating the torso at a machine is contraindicated for people who have low back problems or who have already had a herniated disc because they risk aggravating or reviving this lumbar pathology.



VARIATION SEATED AT A MACHINE



DEEP MUSCLES OF THE ABDOMEN



Rectus abdominis

Gluteus medius

Tensor fasciae latae

Pyramidalis

Pubic symphysis

Quadriceps, rectus femoris

Fascia lata, iliotibial tract

External oblique

Anterior superior iliac spine

Internal oblique, under the aponeurosis

Iliopsoas

Pectineus

Sartorius

Adductor longus

Gracilis

Quadriceps, vastus medialis

Quadriceps, vastus lateralis

Stand on the swivel plate and grasp the handle:

- Rotate the pelvis to one side then to the other, keeping the shoulders fixed. The knees should be slightly bent. Control the rotations.

This exercise mainly works the external and internal obliques and, to a lesser degree, the rectus abdominis.

To feel the effort more intensely on the external and internal obliques, round the back slightly.

Best results are obtained with very long sets.




STRETCHING THE ABDOMINALS

Lie on your belly, resting on your hands with arms extended:


- Slowly raise the torso, tilting slightly back with your head.
- Maintain the position for a few moments, breathing slowly to really feel the stretch of the anterior part of the abdominal girdle.

Variations: You can stretch the abdominal muscles with your hands resting on a bench or your feet on the floor, or you can stretch backward on a Swiss ball.

Comment: Stretching the abdominal girdle is recommended in certain sports, such as throwing events in track and field, especially with the javelin, where good flexibility and good abdominal amplitude are essential for performing the movement perfectly.



COMPARING THE PELVIC TILT IN WOMEN AND MEN




TILT OF THE PELVIS

A woman's pelvis generally tilts more anteriorly than a man's. This anterior tilt pushes the gluteals back more and pulls the pubis in more between the thighs, which gives the impression that the lower belly is pushed out slightly. This small, typically feminine belly contrasts with the vertical abdominal wall that is more commonly seen in men, where the pelvis is tilted forward to a lesser degree.


The special position of the female pelvis prevents a fetus from excessively compressing the viscera during pregnancy as part of its weight presses against the abdominal wall.


A: Anterior superior iliac spine
B: Pubic tubercle


SAGITTAL SECTION OF THE ABDOMEN OF A PREGNANT WOMAN




Comment: The anterior tilt position (anteversion) of the pelvis in a woman allows part of the weight of the fetus to press against the abdominal wall. The muscles of the abdominal wall can be compared to a hammock.









Images have been losslessly embedded. Information about the original file can be found in PDF attachments. Some stats (more in the PDF attachments):

```
{
  "filename": "NDA3NTAzNTAuemlw",
  "filename_decoded": "40750350.zip",
  "filesize": 39832502,
  "md5": "f6e47612d0eab38afeed960d453aa64b",
  "header_md5": "f3195e327e7ad55702438beab227d395",
  "sha1": "6cab10df4f04539f2217327e8fdf478579ec5d0d",
  "sha256": "1145c6f71acd971521eae12fc9395688dcc2bb5b260745c2665295e2d690142d",
  "crc32": 3043651053,
  "zip_password": "",
  "uncompressed_size": 48887816,
  "pdg_dir_name": "STRENGTH TRAINING ANATOMY THIRD EDITION_40750350",
  "pdg_main_pages_found": 192,
  "pdg_main_pages_max": 192,
  "total_pages": 197,
  "total_pixels": 1257789840,
  "pdf_generation_missing_pages": false
}
```